

ADOPTED: 21 April 2016

doi: 10.2903/j.efsa.2016.4484

## Dietary Reference Values for choline

### EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA)

#### Abstract

Following a request from the European Commission, the EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) derives Dietary Reference Values (DRVs) for choline. In this Opinion, the Panel considers dietary choline including choline compounds (e.g. glycerophosphocholine, phosphocholine, phosphatidylcholine, sphingomyelin). The Panel considers that none of the biomarkers of choline intake or status is suitable to derive DRVs for choline. The Panel considers that Average Requirements and Population Reference Intakes for choline cannot be derived for adults, infants and children, and therefore defines Adequate Intakes (AIs). For all adults, the Panel sets an AI at 400 mg/day based on the average observed choline intake in healthy populations in the European Union and in consideration of the amounts of choline needed to replete about 70% of depleted subjects who showed signs of organ dysfunction in a depletion/repletion study. For all infants aged 7–11 months, the Panel proposes an AI of 160 mg/day, based on upwards extrapolation from the estimated choline intake of exclusively breast-fed infants from birth to 6 months. For all children aged 1–17 years, the Panel proposes AIs, based on downwards extrapolation from the adult AI, applying growth factors. These AIs range from 140 mg/day (1–3 years) to 400 mg/day (15–17 years). For pregnant women, the Panel derives an AI of 480 mg/day, calculated by extrapolation from the AI for non-pregnant women and the mean gestational increase in body weight. For lactating women, the amount of choline secreted per day in human milk during the first 6 months of exclusive breastfeeding (120 mg/day) is added to the AI for non-lactating women and an AI of 520 mg/day is set.

© 2016 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

**Keywords:** choline, phosphatidylcholine, observed intake, depletion/repletion study, Adequate Intake, Dietary Reference Value

**Requestor:** European Commission

**Question number:** EFSA-Q-2011-01208

**Correspondence:** [nda@efsa.europa.eu](mailto:nda@efsa.europa.eu)

**Panel members:** Jean-Louis Bresson, Barbara Burlingame, Tara Dean, Susan Fairweather-Tait, Marina Heinonen, Karen-Ildico Hirsch-Ernst, Inge Mangelsdorf, Harry McArdle, Androniki Naska, Monika Neuhäuser-Berthold, Grażyna Nowicka, Kristina Pentieva, Yolanda Sanz, Alfonso Siani, Anders Sjödin, Martin Stern, Daniel Tomé, Dominique Turck, Henk Van Loveren, Marco Vinceti and Peter Willatts

**Acknowledgements:** The Panel wishes to thank the members of the Working Group on Dietary Reference Values for vitamins: Christel Lamberg-Allardt, Monika Neuhäuser-Berthold, Grażyna Nowicka, Kristina Pentieva, Hildegard Przyrembel, Inge Tetens, Daniel Tomé and Dominique Turck for the preparatory work on this scientific opinion and EFSA staff members Fanny Héraud and Francisca Vennemann for the support provided to this scientific opinion.

**Suggested citation:** EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2016. Scientific opinion on Dietary Reference Values for choline. *EFSA Journal* 2016;14(8):4484, 70 pp. doi:10.2903/j.efsa.2016.4484

**ISSN:** 1831-4732

© 2016 European Food Safety Authority. *EFSA Journal* published by John Wiley and Sons Ltd on behalf of European Food Safety Authority.

This is an open access article under the terms of the [Creative Commons Attribution-NoDerivs License](#), which permits use and distribution in any medium, provided the original work is properly cited and no modifications or adaptations are made.



The EFSA Journal is a publication of the European Food Safety Authority, an agency of the European Union.



## Summary

Following a request from the European Commission, the EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA) was asked to deliver a Scientific Opinion on Dietary Reference Values (DRVs) for the European population, including choline.

Choline is a quaternary amine (2-hydroxyethyl-*N,N,N*-trimethylammonium) present in food in free and esterified forms. The main forms present in foods are phosphatidylcholine (PC, lecithin), which is also the main form present in animal tissues, free choline, phosphocholine (PChol), glycerophosphocholine (GPC) and sphingomyelin (SPM), and minor amounts of cytidine-5-diphosphate-choline (CDP-choline) and acetylcholine. Choline, PChol and GPC are water-soluble choline compounds, whereas PC and SPM are lipid-soluble compounds.

Although choline can be synthesised *de novo* by the human body, this synthesis may become insufficient, making choline an essential component of the diet. Choline is predominantly provided via the diet. The human body can form choline either *de novo* by methylation of phosphatidylethanolamine (PE) via the hepatic phosphatidylethanolamine *N*-methyltransferase (PEMT) pathway, or by hydrolysis of PC formed in the CDP-choline pathway in all cells of the body. The PC formed in the PEMT pathway contains substantial amounts of long-chain polyunsaturated fatty acids, like docosahexaenoic acid and arachidonic acid. Both pathways can be stimulated by dietary choline and the PEMT pathway is sensitive to the presence of oestrogens. Choline is an integral part of some phospholipids, which play an important role in the structure and function of membranes. Choline (as PC) plays an important role in the metabolism and transport of lipids and cholesterol by lipoproteins, and is needed for the assembly and secretion of very low-density lipoproteins by the liver. Choline is a precursor of the neurotransmitter acetylcholine, and of betaine, an osmoregulator to which choline is irreversibly oxidised in the liver and kidney. Via betaine, choline is involved in the folate-dependent one-carbon metabolism. Dietary deficiency of choline can cause fatty liver or hepatic steatosis that can result in non-alcoholic fatty liver disease (NAFLD), and can cause liver and muscle damage. This shows that *de novo* production can be insufficient.

Dietary free choline is quickly taken up by a carrier-mediated saturable transport system. PC and GPC from the diet or secreted in the bile, and dietary SPM are hydrolysed by phospholipases (PLs) to liberate choline. Choline and water-soluble choline compounds (PChol and GPC) are rapidly absorbed and appear in plasma predominantly as free choline. Phospholipids (PC and SPM) that have escaped PLs enter the lymph incorporated into chylomicrons. The available data do not allow defining the percentage of intestinal absorption of choline in humans, and the total amount of choline in the human body. Non-absorbed choline is a precursor of trimethylamine (TMA) produced in the gut by anaerobic symbiotic microbes. TMA is efficiently absorbed from the gastrointestinal tract and then converted in the liver to trimethylamine-*N*-oxide (TMAO). Both TMA and TMAO (i.e. total trimethylamine (TTMA)) are eliminated in the urine. Urinary excretion of choline is low in relation to usual dietary intakes, while no human data are available on faecal excretion of choline or choline compounds in relation to dietary intake. Breast milk mainly contains PChol and GPC, besides free choline, PC and SPM, in concentrations depending on the progress of lactation, maternal diet and genotype.

The Panel reviewed possible biomarkers of choline intake and/or status. The Panel considers that the available data do not allow conclusions to be drawn on a dose–response relationship between choline intake or status and plasma choline concentration, and that plasma choline concentrations cannot be used to set DRVs for dietary choline. Plasma concentrations of PC, betaine, dimethylglycine, total homocysteine or TMAO, erythrocyte PC concentration, or urinary betaine and TTMA urinary excretion also cannot be used to set DRVs for dietary choline. The Panel also notes that single-nucleotide polymorphisms in genes coding for enzymes involved in choline metabolism, some of them present with high frequency in the population, can influence the dietary requirement for choline and determine the susceptibility to dietary choline deficiency, but data are insufficient to predict variations in individual choline requirements based on genetic polymorphisms. The Panel considers that the available data on choline intake and health consequences (NAFLD, cardiovascular disease, cancer, birth defects, cognition) cannot be used to set DRVs for dietary choline.

The Panel considers that Average Requirements and Population Reference Intakes for choline cannot be derived for adults, infants and children, and therefore defines Adequate Intakes (AIs).

Dietary total choline intake was calculated based on individual food consumption data that were available to the European Food Safety Authority (EFSA) and classified according to EFSA's food classification system, from healthy populations investigated in 12 national surveys undertaken in nine countries of the European Union (EU), between 2000 and 2011. In the absence of food composition

data on choline in Europe, composition data on free choline and choline compounds from the US Department of Agriculture were used. The total choline intake mean estimates ranged from 75 to 127 mg/day in infants, from 151 to 210 mg/day in children aged from 1 to < 3 years, from 177 to 304 mg/day in children aged from 3 to < 10 years, and from 244 to 373 mg/day among children aged from 10 to < 18 years. The total choline intake mean estimate was 336 mg/day in pregnant adolescents and 356 mg/day in pregnant women. The total choline intake mean estimates ranged from 269 to 444 mg/day and from 332 to 468 mg/day in women and men, respectively, i.e. for all adults: 269–468 mg/day.

The Panel reviewed 11 choline depletion/repletion studies with similar design. Only one reported the amounts of choline needed to replete depleted subjects who showed signs of organ dysfunction. The Panel concludes that choline depletion/repletion studies do not provide sufficient data to calculate Average Requirements for choline, but may be used to inform data on observed choline intakes to set AIs for choline.

For all adults, the Panel set an AI of 400 mg/day. This is based on the midpoint of the range of observed mean intakes in healthy populations in the EU (about 370 mg/day), and in consideration of the results of a depletion/repletion study in which about 70% of the depleted subjects who had developed signs of organ dysfunction were repleted with an intake of about 400 mg/70-kg body weight (bw) per day. Although premenopausal women may have a lower requirement for dietary choline (than postmenopausal women or men) in connection with a potential stimulation of the PEMT pathway by oestrogens, and ranges of estimated mean total choline intake in Europe are slightly lower in women than men, the Panel considered it unnecessary to give sex-specific AIs for adults.

For infants aged 7–11 months, the Panel set an AI of 160 mg/day, based on the estimated intake of choline of exclusively breast-fed infants from birth to 6 months, and upwards extrapolation by allometric scaling, taking into account the difference in reference body weight.

For all children aged 1–17 years, no data are available that would justify different AIs for boys and girls. The Panel set AIs ranging from 140 mg/day (1–3 years) to 400 mg/day (15–17 years). These were set by downwards extrapolation from the adult AI, by allometric scaling, taking into account the difference in reference body weight and applying growth factors. These AIs are supported by total choline intake mean estimates in the EU.

The Panel considered that, although the available intervention studies on choline supplementation in the second half of pregnancy or in lactating women indicate that pregnant or lactating women may need more choline than non-pregnant non-lactating women, the data are not sufficient to allow an estimate of the additional requirement for dietary choline in pregnant or lactating women (above that of non-pregnant non-lactating women).

For pregnant women, the Panel set an AI of 480 mg/day, calculated by isometric scaling from the AI for non-pregnant women, using the mean gestational increase in body weight. For lactating women, the AI for non-lactating women is increased to account for the secretion of choline through breast milk. The Panel set an AI of 520 mg/day, considering an average concentration of choline in mature breast milk of 145 mg/L, and a mean milk transfer during the first 6 months of lactation in exclusively breastfeeding women (0.8 L/day).

## Table of contents

Abstract.....	1
Summary.....	3
Background as provided by the European Commission.....	7
Terms of reference as provided by the European Commission.....	7
Assessment.....	8
1. Introduction.....	8
2. Definition/category.....	8
2.1. Chemistry.....	8
2.2. Function of choline.....	9
2.2.1. Biochemical functions.....	9
2.2.2. Health consequences of deficiency and excess.....	9
2.2.2.1. Deficiency.....	9
2.2.2.2. Excess.....	10
2.3. Physiology and metabolism.....	10
2.3.1. Intestinal absorption.....	10
2.3.2. Transport in blood.....	11
2.3.3. Distribution to tissues.....	11
2.3.4. Storage.....	11
2.3.5. Metabolism.....	12
2.3.5.1. Metabolism of choline and synthesis of phosphatidylcholine (PC).....	12
2.3.5.2. Degradation.....	13
2.3.5.2.1. Choline oxidation to betaine.....	14
2.3.5.2.2. Microbial choline degradation to trimethylamine (TMA).....	14
2.3.6. Elimination.....	14
2.3.6.1. Urine.....	14
2.3.6.1.1. Choline and trimethylamine- <i>N</i> -oxide (TMAO).....	15
2.3.6.1.2. Betaine and dimethylglycine (DMG).....	15
2.3.6.1.3. Conclusion on urinary excretion.....	16
2.3.6.2. Faeces.....	16
2.3.6.3. Breast milk.....	16
2.3.7. Interaction with other nutrients: folate.....	17
2.4. Biomarkers.....	18
2.4.1. Plasma/serum concentration of free choline and choline-compounds and metabolites.....	18
2.4.1.1. Adults.....	18
2.4.1.2. Pregnancy and lactation.....	19
2.4.1.3. Infants.....	20
2.4.1.4. Conclusion on plasma/serum concentration of choline and choline-compounds and metabolites.....	20
2.4.2. Total trimethylamine (TTMA) hepatic production.....	20
2.4.3. Plasma total homocysteine.....	20
2.4.4. Urinary betaine excretion.....	21
2.4.5. Conclusions on biomarkers.....	21
2.5. Effects of genotypes involved in choline metabolism.....	21
2.5.1. Influence of polymorphisms in pregnancy and lactation.....	22
2.5.2. Conclusion on effects of genotypes.....	22
3. Dietary sources and intake data.....	23
3.1. Dietary sources.....	23
3.2. Dietary intake.....	23
3.2.1. Dietary intake in EU countries.....	23
3.2.2. Dietary intake in non-EU countries.....	24
3.2.3. Conclusion on dietary intake.....	24
4. Overview of dietary reference values and recommendations.....	24
4.1. Adults.....	25
4.2. Infants and children.....	25
4.3. Pregnancy.....	25
4.4. Lactation.....	25
5. Criteria (endpoints) on which to base Dietary Reference Values.....	26
5.1. Indicators of choline requirement.....	26
5.1.1. Adults.....	26
5.1.1.1. Study goals.....	26

5.1.1.2.	Study design.....	27
5.1.1.3.	Number of subjects and choline intake.....	27
5.1.1.4.	Summary.....	29
5.1.2.	Infants and children.....	29
5.1.3.	Pregnancy and lactation.....	29
5.1.3.1.	Effect of total intake of choline in pregnant (vs non-pregnant) women and the offspring.....	29
5.1.3.2.	Effect of total intake of choline in pregnant (vs non-pregnant) women on the dynamics of choline-related metabolic pathways.....	30
5.1.3.3.	<i>Ex-vivo</i> studies in placental samples.....	31
5.1.3.4.	Effect of total intake of choline on maternal plasma and breast milk during lactation.....	31
5.1.3.5.	Conclusion on pregnancy and lactation.....	31
5.2.	Choline intake and health consequences.....	32
5.2.1.	Non-alcoholic fatty liver disease.....	33
5.2.2.	Cardiovascular disease.....	33
5.2.3.	Cancer.....	33
5.2.3.1.	Colon/rectum.....	33
5.2.3.2.	Breast cancer.....	34
5.2.3.3.	Other cancers (oesophageal, prostate and ovarian cancers).....	34
5.2.3.4.	Conclusions.....	35
5.2.4.	Neural tube defects.....	35
5.2.5.	Cognition.....	35
5.2.6.	Conclusion on choline intake and health consequences.....	36
6.	Data on which to base dietary reference values.....	36
6.1.	Adults.....	36
6.2.	Infants.....	37
6.3.	Children.....	38
6.4.	Pregnancy.....	39
6.5.	Lactation.....	40
	Conclusions.....	40
	Recommendations for research.....	40
	References.....	41
	Abbreviations.....	48
	Appendix A – Concentrations of free and total choline in breast milk of healthy lactating mothers.....	51
	Appendix B – Intervention and observational studies on the relationship between dietary choline and plasma homocysteine concentration.....	56
	Appendix C – SNPs of genes coding for enzymes involved in choline metabolism and their impact on choline requirement and/or risk to develop organ dysfunction while being fed a low-choline diet.....	59
	Appendix D – Depletion/repletion studies for choline.....	62



## Background as provided by the European Commission

The scientific advice on nutrient intakes is important as the basis of Community action in the field of nutrition, for example such advice has in the past been used as the basis of nutrition labelling. The Scientific Committee for Food (SCF 1993) report on nutrient and energy intakes for the European Community dates from 1993. There is a need to review and if necessary to update these earlier recommendations to ensure that the Community action in the area of nutrition is underpinned by the latest scientific advice.

In 1993, the SCF adopted an opinion on the nutrient and energy intakes for the European Community.<sup>1</sup> The report provided Reference Intakes for energy, certain macronutrients and micronutrients, but it did not include certain substances of physiological importance, for example dietary fibre.

Since then new scientific data have become available for some of the nutrients, and scientific advisory bodies in many European Union (EU) Member States and in the United States have reported on recommended dietary intakes. For a number of nutrients, these newly established (national) recommendations differ from the reference intakes in the SCF (1993) report. Although there is considerable consensus between these newly derived (national) recommendations, differing opinions remain on some of the recommendations. Therefore, there is a need to review the existing EU Reference Intakes in the light of new scientific evidence, and taking into account the more recently reported national recommendations. There is also a need to include dietary components that were not covered in the SCF opinion of 1993, such as dietary fibre, and to consider whether it might be appropriate to establish reference intakes for other (essential) substances with a physiological effect.

In this context, EFSA is requested to consider the existing Population Reference Intakes for energy, micro- and macronutrients and certain other dietary components, to review and complete the SCF recommendations, in the light of new evidence, and in addition advise on a Population Reference Intakes for dietary fibre.

For communication of nutrition and healthy eating messages to the public, it is generally more appropriate to express recommendations for the intake of individual nutrients or substances in food-based terms. In this context, EFSA is asked to provide assistance on the translation of nutrient-based recommendations for a healthy diet into food-based recommendations intended for the population as a whole.

## Terms of reference as provided by the European Commission

In accordance with Article 29 (1)(a) and Article 31 of Regulation (EC) No 178/2002<sup>2</sup>, the Commission requests EFSA to review the existing advice of the Scientific Committee for Food on Population Reference Intakes for energy, nutrients and other substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

In the first instance, EFSA is asked to provide advice on energy, macronutrients and dietary fibre. Specifically, advice is requested on the following dietary components:

- Carbohydrates, including sugars;
- Fats, including saturated fatty acids, polyunsaturated fatty acids and monounsaturated fatty acids, *trans* fatty acids;
- Protein;
- Dietary fibre.

Following on from the first part of the task, EFSA is asked to advise on population reference intakes of micronutrients in the diet and, if considered appropriate, other essential substances with a nutritional or physiological effect in the context of a balanced diet which, when part of an overall healthy lifestyle, contribute to good health through optimal nutrition.

Finally, EFSA is asked to provide guidance on the translation of nutrient-based dietary advice into guidance, intended for the European population as a whole, on the contribution of different foods or categories of foods to an overall diet that would help to maintain good health through optimal nutrition (food-based dietary guidelines).

<sup>1</sup> Scientific Committee for Food, Nutrient and energy intakes for the European Community, Reports of the Scientific Committee for Food 31st series, Office for Official Publication of the European Communities, Luxembourg, 1993.

<sup>2</sup> Regulation (EC) No 178/2002 of the European Parliament and of the Council of 28 January 2002 laying down the general principles and requirements of food law, establishing the European Food Safety Authority and laying down procedures in matters of food safety. OJ L 31, 1.2.2002, p. 1–24.

## Assessment

### 1. Introduction

Choline is a water-soluble organic compound needed for normal functioning of the body. Although choline can be synthesised *de novo* by the human body, this synthesis may become insufficient, making choline an essential component of the diet (Ueland, 2011).


In 1993, the SCF adopted an opinion on nutrient and energy intakes for the European Community and considered that there was no evidence for the necessity of an intake of choline via the diet for persons older than 6 months (SCF, 1993). As it was unclear if young infants depend on exogenous sources of choline and because choline is an integral component of human milk, the addition of choline to infant formula with a minimum level of 7 mg of choline/100 kcal was made mandatory.<sup>3</sup>

The purpose of this Opinion is to review the available evidence to assess whether it might inform the setting of Dietary Reference Values (DRVs) for choline. The Panel focuses in this Scientific Opinion on dietary choline including choline containing compounds.

### 2. Definition/category

#### 2.1. Chemistry

Choline, 2-hydroxyethyl-*N,N,N*-trimethylammonium (2-hydroxy-*N,N,N*-trimethylethanammonium, IUPAC, molar mass 104.17 g/mol) is a quaternary amine. In foods, it is present as free choline and in esterified forms, mainly as phosphatidylcholine (PC, lecithin), phosphocholine (PChol), glycerophosphocholine (GPC) and sphingomyelin (SPM) (Figure 1), and minor amounts of cytidine-5-diphosphate-choline (CDP-choline) and acetylcholine (Ueland, 2011). PC accounts for approximately 95% of total choline found in animal tissues. Choline, PChol and GPC are water-soluble choline compounds, whereas PC and SPM are lipid soluble.



**Figure 1:** Chemical formulas of choline, glycerophosphocholine, phosphocholine, phosphatidylcholine and sphingomyelin

Choline is a component of some phospholipids. Phospholipids are derived from either glycerol or sphingosine, an amino alcohol with a long unsaturated hydrocarbon chain (C18). Phosphoglycerides consist of a glycerol of which the hydroxyl groups at C1 and C2 are esterified to the carboxyl groups of two fatty acids, while the hydroxyl group at C3 is esterified to PChol (or other phosphorylated alcohols derived from ethanolamine, serine or inositol). SPM consists of sphingosine, which amino group is linked to a fatty acid by an amide bond and which primary alcohol group is esterified to PChol.

<sup>3</sup> Commission Directive 2006/141/EC of 22 December 2006 on infant formulae and follow-on formulae and amending Directive 1999/21/EC, OJ L 401, 30.12.2006, p. 1.



## 2.2. Function of choline

### 2.2.1. Biochemical functions

Choline has a number of important functions: it is a precursor for the phospholipid PC (Section 2.1); it is involved in the metabolism and transport of lipids and in the folate-dependent one-carbon metabolism; and it is a precursor of acetylcholine and of betaine.

Choline is an integral part of some phospholipids (Section 2.1). Phospholipids are abundant in all biological membranes (40–50% of phospholipids of cellular membranes consist of PC (Zeisel, 2006)), where they play an important role in the structure and function of membranes, including signalling and transport, and they are also a constituent of the surfactant complex in the lung (Dushianthan et al., 2014).

Choline plays an important role in the metabolism and transport of lipids and cholesterol. PC makes up 70–95% of phospholipids in lipoproteins (Zeisel, 2006) and is needed for normal assembly and secretion of very low-density lipoproteins (VLDL) in the liver (Vance et al., 2007).

Choline is acetylated in cholinergic neurons to form acetylcholine, a key neurotransmitter involved in functions like memory storage and muscle control (IOM, 1998; Ueland, 2011). Pre- and postnatal choline availability has been shown to be important for neurodevelopment in animals (Meck and Williams, 2003).

In the liver and kidney, choline is irreversibly oxidised, by a mitochondrial choline oxidase (also called choline dehydrogenase (CHDH)) and betaine aldehyde dehydrogenase (BADH), to betaine (Lin and Wu, 1986) (Sections 2.3.5.2.1 and 2.3.5.2.2). Betaine serves as an osmoregulator and is a substrate in the betaine-homocysteine methyltransferase (BHMT) reaction. This reaction links choline and betaine to the folate-dependent one-carbon metabolism (Figure 2, Sections 2.3.5 and 2.3.7). Choline and betaine are important sources of one-carbon units, in particular during folate deficiency (Ueland, 2011). In remethylating homocysteine (Hcy) to methionine, choline contributes, via betaine, to the availability of *S*-adenosyl-methionine (SAM) as the universal methyl-group donor (Figure 2, Section 2.3.5). For example, the methyl group of SAM can be transferred to cytosine residues adjacent to guanine (CpG) of DNA or to histones at specific lysine sites, thereby contributing to epigenetic modification and potentially exerting effects on gene expression (Mehedint and Zeisel, 2013).

### 2.2.2. Health consequences of deficiency and excess

#### 2.2.2.1. Deficiency

Dietary deficiency of choline can cause fatty liver (hepatic steatosis, which can result in non-alcoholic fatty liver disease (NAFLD)) (Buchman et al., 1995), and liver (Zeisel et al., 1991) and muscle damage as indicated by an increase in creatine phosphokinase (CK) concentration in serum (Fischer et al., 2007). Hepatic steatosis may be due to impaired triacylglycerol (TAG) transport out of the liver. As PC is an essential component of VLDL, the lipoprotein responsible for transporting TAG out of the liver (Section 2.2.1), TAG cannot be exported in case of choline deficiency and accumulates in the hepatocytes (Cole et al., 2012). Hepatic steatosis can progress to liver damage with release of liver enzymes into the blood. This release of enzymes from the liver into the blood may follow induction of apoptosis and cell membrane fragility (da Costa et al., 2006a; Fischer et al., 2007). In serum of 41 long-term parenterally fed subjects, both alanine aminotransferase (ALT) and aspartate aminotransferase (AST) concentrations were significantly and negatively associated with the concentration of free choline ( $r = -0.34$ ,  $p = 0.03$ ;  $r = -0.37$ ,  $p = 0.02$ , respectively), but not with that of phospholipid-bound choline (Buchman et al., 1993). In this study, the concentration of free choline in serum was low, i.e. one-third of the reference values used by the authors, while that of PC was normal.

The susceptibility to develop NAFLD was found to be related to polymorphisms of the gene for phosphatidylethanolamine *N*-methyltransferase (PEMT) (Song et al., 2005) with loss of oestrogen receptor binding (Resseguie et al., 2007, 2011), as well as to polymorphisms of other enzymes involved in choline metabolism (CHDH and 5,10-methylenetetrahydrofolate dehydrogenase 1 (MTHFD1)) (Section 2.5 and Appendix C). Premenopausal women developed signs of choline deficiency less commonly than postmenopausal women or men, possibly as a consequence of upregulation of hepatic PEMT by oestrogen, leading to an increase in the endogenous synthesis of PC (Fischer et al., 2007; Zeisel, 2007). PEMT is important for this endogenous synthesis of PC in case of insufficient dietary choline intake (Figure 2, Section 2.3.5). The amount of dietary choline to prevent organ damage or to maintain normal organ function varies between people (Section 5.1.2). In addition, there is some evidence that

the susceptibility to develop fatty liver with choline deficiency is influenced by the gastrointestinal microbiome (Spencer et al., 2011) (Appendix D).

Zeisel (2012) reviewed the potential effects of choline deficiency on gene expression via epigenetic marks and DNA integrity that could result in increased mutation rates and thereby increased risks of certain cancers. An influence on the risk of breast cancer of single nucleotide polymorphisms (SNPs) of several genes involved in choline metabolism and enhancing the requirement for dietary choline has been observed in large epidemiological studies (Xu et al., 2008, 2009) (Appendix C and Section 2.5).

In subjects that received a choline diet providing < 50 mg choline/70-kg body weight (bw) per day, fasting plasma concentration of total homocysteine (tHcy) significantly increased among those with clinical expression of choline deficiency compared with baseline (da Costa et al., 2005; Fischer et al., 2007) (Section 5.1.1 and Appendix D). However, many factors besides dietary or endogenous choline determine tHcy concentration in plasma (Section 2.4.3) (EFSA NDA Panel, 2014b, 2015).

#### 2.2.2.2. Excess

The SCF did not consider choline when setting Tolerable Upper Intake Levels (ULs) for vitamins and minerals. The US Institute of Medicine (IOM, 1998) defined a UL for adults based on a study in seven patients with Alzheimer's dementia, where the oral administration of 7.5 g/day of choline (as chloride) had a hypotensive effect accompanied by nausea and diarrhoea (Boyd et al., 1977). Similar gastrointestinal effects and a fishy body odour were observed in therapeutic studies with choline (8–20 g/day) on individuals with tardive dyskinesia and Huntington's disease (Growdon et al., 1977; Gelenberg et al., 1979; Lawrence et al., 1980). The IOM considered 7.5 g/day of choline as the Lowest Observed Adverse Effect Level (LOAEL), and after the application of an uncertainty factor of 2 and rounding, set a UL of 3.5 g choline/day for adults. No UL was established for infants and ULs for children were derived from the adult value by allometric scaling (exponent 0.75) according to reference body weights.

An association between an increased risk of cardiovascular diseases (CVD) and 'higher intake' of choline, which possibly exceeds the intestinal absorption capacity for dietary free choline, has been suggested by a metabolomic study (Wang et al., 2011), which investigated the relationship between plasma choline and trimethylamine-*N*-oxide (TMAO) concentrations and risk of CVD. Non-absorbed choline will become available to microbial degradation, predominantly to trimethylamine (TMA), which is metabolised in the liver to TMAO (Sections 2.3.1 and 2.3.5.2.2). TMA has been found to promote atherosclerosis in animals (Wang et al., 2011, 2014; Bennett et al., 2013; Tang et al., 2013). TMA has also been suggested to be involved in depression, neurological symptoms, teratogenic effects in humans as well as in the potential formation of the carcinogen *N*-nitrosodimethylamine (for a review, see Bain et al. (2005)). These are indirect adverse effects of choline, depending both on a 'high' dietary amount and a specific gut microbiome (Wang et al., 2011). However, the dietary intake of choline was not reported in these studies.

### 2.3. Physiology and metabolism

#### 2.3.1. Intestinal absorption

Dietary free choline is quickly taken up by the enterocytes, mediated by the saturable organic cation transporters (OCTs) (choline transporter-like protein 1 (CTL1) or solute carrier family 44 member 1 (SLC44A1); Section 2.3.3), which rely on facilitated diffusion governed by the choline concentration gradient and the electrical potential across the membrane, then free choline is cleared from the plasma within about 3 h (Zeisel et al., 1980; Jope et al., 1982). Dietary PC increases plasma free choline concentration for 8–12 h, without a significant rise in PC concentration in plasma (Zeisel et al., 1980; Jope et al., 1982). PChol and GPC are rapidly absorbed and appear in plasma predominantly as free choline.

PC and GPC from the diet or secreted in the bile are hydrolysed by phospholipases (PLs) to liberate choline (Zeisel and Blusztajn, 1994). Water-soluble choline compounds (PChol and GPC) can also enter the portal circulation of the liver intact. Lipid-soluble compounds (PC and SPM) are either hydrolysed by PLs or enter the lymph incorporated into chylomicrons.

Unabsorbed choline is catabolised by the intestinal microbiota to TMA (Sections 2.2.2.2 and 2.3.5.2.2). TMA is absorbed from the gastrointestinal tract and converted to TMAO in the liver.

The Panel notes that the amount of choline absorbed is restricted by the capacity of the transport system via the saturable CTL1 or SLC44A1. The Panel notes that the available data do not allow defining the percentage of intestinal absorption of choline in humans.

### 2.3.2. Transport in blood

Free choline is transported in the aqueous phase of plasma, whereas phosphorylated choline compounds (i.e. PC, PChol, GPC, SPM) are associated with or are part of lipoproteins.

### 2.3.3. Distribution to tissues

As free choline is a charged hydrophilic cation, it needs transport mechanisms to cross biological membranes. Three transport mechanisms are known (Fagone and Jackowski, 2013).

The first is a sodium- and chloride-dependent high-affinity ( $K_m < 10 \mu\text{M}$ ) (Okuda and Haga, 2000) carrier-mediated saturable uptake system in presynaptic cholinergic nerve terminals, that is linked to acetylcholine synthesis (Section 2.2.1). The transporter is the high-affinity choline transporter (CHT; solute carrier family 5 member 7 encoded by *SLC5A7*) that needs adenosine triphosphate (ATP) hydrolysis. Disturbing the integrity of the cell membrane can reduce choline availability for acetylcholine synthesis and diminish cholinergic transmission (Cuddy et al., 2014).

The second transport mechanism is a sodium-independent low-affinity carrier-mediated saturable mechanism (CTL1 or *SLC44A1*) in all tissues. This mechanism is energised by ATP hydrolysis, with an average affinity ( $K_m$ ) for choline of  $> 20\text{--}200 \mu\text{M}$ . It is present in enterocytes, hepatocytes, kidneys, placental tissue, mitochondria and synaptosomes, and supplies choline for the synthesis of PC and SPM as well as of betaine (Sections 2.2.1 and 2.3.5). This uptake is stereospecific and can be inhibited by similar nitrogen-methyl compounds and by high concentrations of choline (Michel and Bakovic, 2012).

The third transport mechanism is a sodium-independent saturable uptake mechanism (a member of the solute carrier 22 family), for choline to cross the blood–brain barrier and erythrocyte membranes by facilitated diffusion. Its affinity to choline is similar to the high-affinity mechanism, but it is not linked to acetylcholine synthesis (Cornford et al., 1980; Lockman and Allen, 2002).

Choline uptake by the mammary epithelium occurs by an energy-dependent saturable transport system, but with higher maternal choline supply non-saturable transport can also occur. Choline is metabolised within the mammary epithelium to PChol and other choline compounds or to a lesser extent via degradation pathways (Fischer et al., 2010b; Davenport et al., 2015) (Sections 2.3.6, 2.4.1.2 and 5.1.3). The size of the efflux of choline compounds from the mammary epithelium occurs via exocytosis or as a component of the milk fat globule (Davenport and Caudill, 2013).

Choline crosses the placenta via a specific transport system on both the maternal and fetal side of the syncytiotrophoblast (Baumgartner et al., 2015), with an apparent small excess (about 4%) preferential towards the fetal circulation, as demonstrated in perfusion studies with [ $^3\text{H}$ ]-choline (Sweiry et al., 1986). Umbilical cord blood free choline concentration is about three times that of maternal blood (Visentin et al., 2015) (Section 2.4.1.2).

### 2.3.4. Storage

Choline is stored in tissues either as membrane-bound phospholipids or as intracellular PC or GPC (Zeisel and Blusztajn, 1994). Choline is stored in the brain as membrane-bound phospholipids, which are hydrolysed by choline acetyltransferase to provide choline for acetylcholine synthesis (Section 2.2.1). In most animal tissues, PC accounts for 95% of the total choline content, the remaining 5% are choline, PChol, GPC, CDP-choline and acetylcholine (Li and Vance, 2008).

The content of choline and its metabolites in the body is balanced by two pathways of acquisition, i.e. either diet and the CDP pathway, or the PEMT pathway (Sections 2.2.2.1 and 2.3.5, Figure 2), and two pathways of depletion, i.e. either choline oxidation or the secretion of PC in the bile and, to a lesser extent, by the intestinal mucosa (Li and Vance, 2008; Eehalt et al., 2010) (Sections 2.3.5.2.1 and 2.3.6.2). Among bile phospholipids, 95% is PC, of which about 40% returns to the liver. Choline imbalances can be compensated by adaptive increases in PEMT activity, by recycling of choline, decreased oxidation of choline, reabsorption of biliary PC, and by redistribution of tissue choline to maintain homeostasis particularly in the brain and liver (Li et al., 2007; Li and Vance, 2008).

Regarding the choline content of adult tissues, data obtained with proton (hydrogen-1 [ $^1\text{H}$ ]) magnetic resonance spectroscopy (MRS) are available. This method measures, besides choline as such, primarily GPC and PChol, but also includes phosphatidylethanolamine (PE), glycerolPE, betaine, myo-inositol and taurine; however, it does not include all choline lipids in membranes. The choline content of human liver has been measured *in vivo* to be on average 8.6 mmol/kg or 894 mg/kg wet weight (range 3.8–17.6 mmol/kg;  $n = 44$  including 24 women, mean age  $46 \pm 17$  years) using proton MRS (Ouwerkerk et al., 2012). The choline content of quadriceps muscle was in the range 6.7–13 mmol/kg or


697–1,352 mg/kg (n = 7 including 4 women, mean age 37.7 years, range 28–50 years) (Fayad et al., 2010). The choline content in parietal white matter of the brain was (mean ± SD) 1.73 ± 0.24 mmol/L or 180 ± 25 mg/L (n = 20 including 11 women, mean age 29.4 ± 7.4 years) (Mazzetti et al., 2013).

Regarding the fetus, infant and young child, phospholipids in the brain increase twofold in the cortex (and threefold in the white matter) from the 10th week of gestation to the age of 2 years (Svennerholm and Vanier, 1972). This study shows a relative continuous decrease in choline phosphoglycerides from 50% of total phospholipids in the cerebral cortex of the fetus to 45% in infants at term and 38% in children at 2 years of age. In this study, SPM shows a continuous increase, from 3% of total phospholipids in the cerebral cortex of the fetus to 5% in infants at term and 10% in children at 2 years of age.

Regarding the placenta, total lipid content is 14 ± 1.0 mg/g dry tissue at term, and is rich in phospholipids (about 80% of total lipids), of which 42.1 ± 7.3% were choline glycerophospholipids (Bayon et al., 1993; Bitsanis et al., 2005). The long-chain polyunsaturated fatty acids (LC-PUFAs), arachidonic acid (ARA) and docosahexaenoic acid (DHA), are found in high proportion (about 40% of the phospholipid fatty acids) in all phospholipid classes (Bayon et al., 1993; Bitsanis et al., 2005). The placenta is one of the human organs most rich in free choline (14.6 mg/100 g wet weight) and this concentration decreases by 50% in (pre)eclampsia (Mischel, 1956).

The Panel notes that no data are available on the total amount of choline in the human body. The Panel also notes that there is a lack of data on the choline accretion in the fetus and placenta during the duration of pregnancy.

### 2.3.5. Metabolism



Left shows the endogenous synthesis of PC (PEMT pathway); right the synthesis of PC from (dietary) choline (CDP pathway). BADH: betaine aldehyde dehydrogenase; BHMT: betaine homocysteine methyltransferase; CCT: phosphocholine cytidyltransferase; CDP-choline, cytidine diphosphocholine; CHK: choline kinase; CHDH: choline oxidase (or dehydrogenase); CPT: CDP-choline diacylglycerol choline phosphotransferase; DMG: dimethylglycine; Hcy: homocysteine; methyl-THF: methyltetrahydrofolate; MS: methionine synthase; PChol: phosphocholine; PE: phosphatidylethanolamine; PEMT: phosphatidylethanolamine N-methyltransferase; PC: phosphatidylcholine; SAH: S-adenosylhomocysteine; SAH-H: S-adenosylhomocysteine hydrolase; SAM: S-adenosylmethionine; THF: tetrahydrofolate.

**Figure 2:** PC synthesis and choline metabolism and its involvement in folate-dependent one-carbon metabolism

#### 2.3.5.1. Metabolism of choline and synthesis of phosphatidylcholine (PC)

Besides dietary intake, choline in the body can be generated *de novo* via the hepatic PEMT pathway. Choline from both dietary and endogenous sources is incorporated into PC. PC is synthesised in all cells from choline (Li and Vance, 2008).

The predominant pathway of PC synthesis in all cells is via the CDP-choline pathway. Free choline, taken up into cells or generated by hydrolysis of choline compounds (Figure 2, right side), is



phosphorylated by choline kinase (CHK) to PChol or oxidised to betaine in some cell types like liver and kidney. PChol reacts with cytidine triphosphate (CTP) to form cytidine 5-diphosphate choline (CDP-choline) by phosphocholine cytidyltransferase (CCT). CDP-choline is esterified with diacylglycerol by choline phosphotransferase (CPT) or the choline/ethanolaminephosphotransferase (CEPT) to form PC (Li and Vance, 2008).

The other pathway of PC synthesis in the human body *de novo* (PEMT pathway) starts from 3-phosphoglycerate, which receives two acyl groups from acyl-coenzyme A and is converted to a phosphatidate (not shown in Figure 2). Phosphatidate can react with CTP to form cytidine diphosphate-diacylglycerol, whose hydroxyl group can react with serine to form phosphatidylserine that is decarboxylated to PE. PE can then be methylated in the liver to synthesise PC (Figure 2, left side). This reaction is catalysed by PEMT, which is dependent on SAM, and consumes three molecules of SAM, while releasing three molecules of *S*-adenosylhomocysteine (SAH) per molecule of formed PC. Quantitatively, this appears to be the most important SAM-dependent transmethylation reaction and source of Hcy in mammals (Stead et al., 2006). The PEMT pathway is mostly active in the liver, but some low activity has been described in other tissues, e.g. in adrenal medulla, mammary gland and adipose tissue at about 0.1% of the hepatic activity (Vance, 2014).

The PEMT pathway accounts for 30% of hepatic PC synthesis in rodents, while 70% are produced from choline via the CDP-choline pathway (Reo et al., 2002; Li and Vance, 2008). The gene for PEMT has multiple oestrogen-responsive elements and its transcription is enhanced by oestradiol *in vitro*. Oestrogen enhanced activity of PEMT can provide for the increased demand for choline during pregnancy when oestrogen concentrations are high (Resseguie et al., 2007) (Sections 2.2.2.1 and 5.1.3).

PC derived via the two different pathways apparently enters separate pools. PC formed in the hepatic PEMT pathway differs from that generated via the CDP-choline pathway, in that it contains primarily LC-PUFAs like DHA and ARA instead of medium-chain, mono- and bi-unsaturated and saturated fatty acids. This has been demonstrated in studies with deuterated choline and ethanolamine in rat and mouse liver, and in mice and humans after parenteral administration of methyl-D<sub>9</sub>-choline<sup>4</sup> (DeLong et al., 1999; Pynn et al., 2011) and using multiple isotopomer distribution analysis (MIDA) (Pynn et al., 2011). In addition, in 21 healthy non-pregnant women randomised to consume for 12 weeks either 480 or 930 mg choline/day (about 20% of which was provided as methyl-D<sub>9</sub>-choline for the last 6 weeks), Yan et al. (2013) demonstrated that the higher choline intake (930 mg/day) favours the use of the PEMT pathway (relative to CDP-choline pathway), and yielded a significantly higher isotope enrichment in plasma PC-DHA (West et al., 2013) (Section 5.1.3).

The ratio of PC-DHA to total PC in plasma is considered a surrogate measure for hepatic PEMT activity (3% of total plasma PC is PC-DHA). It is significantly greater ( $p < 0.01$ ) in premenopausal women than in men or in postmenopausal women. It is significantly lower ( $p < 0.05$ ) in premenopausal women homozygous for the loss-of-function rs12325817 SNP of the *PEMT* gene than in women with the wild-type of *PEMT*. This has been confirmed by measuring PEMT activity in liver biopsies together with the PC-DHA concentration (da Costa et al., 2011) (Appendix D).

The Panel notes that the PC required by the body can be derived from dietary choline and from endogenous synthesis, but is distributed into different pools and carries different fatty acids. The PC formed in the PEMT pathway contains substantial amounts of LC-PUFAs, like DHA and ARA, while the PC formed in the CDP-choline pathway does not. The PEMT pathway is mostly active in the liver, but some low activity has been described in, e.g. in adrenal medulla, mammary gland and adipose tissue. The CDP-choline pathway is present in all cells of the body. Both can be stimulated by dietary choline. Moreover, the PEMT pathway is sensitive to the presence of oestrogens.

### 2.3.5.2. Degradation

Catabolism of phospholipids is initiated by PLs hydrolysing their respective bonds: i.e. PLA1 and PLA2 hydrolyse fatty-acyl bonds (e.g. PC to lysophosphatidylcholine (lyso-PC)), PLC glycerophosphate bonds, and PLD choline phosphate ester bonds. Further, lysoPL degrades lysophosphatidylcholine, which is subsequently converted to GPC and further hydrolysed to choline by a phosphodiesterase (Lockman and Allen, 2002).

<sup>4</sup> Methyl-D<sub>9</sub>-choline, with fully deuterated methyl groups, can either be converted via the CDP-choline pathway to D<sub>9</sub>-PC or by oxidation to D<sub>9</sub>-betaine that will transfer D<sub>3</sub>-methyl groups to homocysteine via BHMT, forming D<sub>3</sub>-methionine and D<sub>6</sub>-DMG. D<sub>3</sub>-methionine can transfer deuterated methyl groups to PE via PEMT, forming predominantly D<sub>3</sub>-PC and D<sub>6</sub>-PC. By estimating the enrichment of the different metabolites and the ratios of deuterated isotopomers, an assessment of the metabolic fluxes is possible (Pynn et al., 2011).

#### 2.3.5.2.1. Choline oxidation to betaine

Oxidation of choline in the liver and kidney produces, in a two-step enzymatic reaction, first betaine aldehyde by mitochondrial CHDH, and then betaine by mitochondrial or cytoplasmic BADH (Lin and Wu, 1986) (Figure 2). Mitochondrial betaine synthesis from choline is controlled by choline transport across the mitochondrial membrane (O'Donoghue et al., 2009). The formation of betaine links choline to the folate-dependent one-carbon metabolism, because betaine is the methyl-group donor in the BHMT reaction (Sections 2.2.1, 2.3.6.1.2 and 2.3.7). This reaction converts Hcy in the liver and kidney to methionine and releases dimethylglycine (DMG), which is converted into sarcosine and methylene-tetrahydrofolate with tetrahydrofolate (THF) as methyl-group acceptor. The resultant sarcosine can be degraded into glycine or be excreted in the urine, while methylene-THF can be reduced to methyl-THF by methylene-THF reductase (MTHFR) (Ueland et al., 2005) (Section 2.5).

#### 2.3.5.2.2. Microbial choline degradation to trimethylamine (TMA)

Non-absorbed choline is one of the precursors of TMA produced in the gut by anaerobic symbiotic microbes (Zhang et al., 1999; Craciun and Balskus, 2012) (Section 2.2.2.2). TMA is efficiently absorbed from the gastrointestinal tract (Al-Waiz et al., 1987), and then converted in the liver to TMAO by the flavin-containing monooxygenase isoform 3 enzyme (FMO3) (Lang et al., 1998). Both TMA and TMAO are eliminated in the urine (urinary total TMA i.e. TTMA = TMA plus TMAO).

TMA has an unpleasant fishy odour and can result in a corresponding fishy body odour when either choline intake is 'high' (Section 2.2.2.2), the intestinal microbiota is disturbed or the subjects suffer from autosomal-recessive trimethylaminuria due to defects in FMO3 (Mitchell and Smith, 2001; Zeisel et al., 2003).

On 'normal' diets, only milligram amounts of TMA were excreted in the urine of healthy subjects and subjects with liver cirrhosis, but when single choline doses of 2–8 g as bicarbonate were given on separate occasions, about 69% of choline nitrogen was excreted in the urine as TMA nitrogen (De la Hueriga and Popper, 1951).

In a study in six healthy males, measuring the conversion of single oral doses of 15 mmol of choline or PC (i.e. 2.1 and 11.65 g, respectively, given on separate occasions at least 2 weeks apart) into urinary TTMA, about 63% of choline appeared as urinary TTMA within 3 days after ingestion (Zhang et al., 1999). In this study, PC did not lead to similar increases in urinary TTMA concentration (0.5–2% of the administered dose).

However, a double-blind randomised controlled trial (RCT) in six healthy volunteers (four women), consuming single increasing amounts of PC separated by 2–4 weeks (119 up to 714 mg/day of choline, mainly as PC, in the form of egg yolk(s)) in addition to a low-choline diet,<sup>5</sup> demonstrated that an intake of increasing amounts of PC resulted in a rise in TMAO concentrations in both plasma and urine (Miller et al., 2014). TMAO concentration in plasma increased in five of six subjects after egg ingestion, with a peak after 6–8 h; however, there was great interindividual variability. TMAO concentration in urine in the 24 h after egg yolk ingestion increased in proportion to the amount of PC ingested (11–15% of the total ingested choline). The authors also found differences in the profile of the faecal microbiome and in the gene for the FMO3 enzyme (the SNP *FMO3* G566A, rs2266782 is associated with a 25% reduction in the enzyme activity) between the study participants. This may explain the variable responses of plasma and urinary TMAO concentrations to PC intake.

The Panel notes a relationship between dietary choline, microbial metabolism of choline to TMA, hepatic TMAO production and urinary TTMA excretion. The Panel notes as well an influence of other dietary, genetic and environmental factors on TMA production. The Panel concludes that a dose–response relationship between dietary choline and hepatic TTMA production cannot be established.

### 2.3.6. Elimination

#### 2.3.6.1. Urine

The kidneys accumulate choline via the sodium-independent low-affinity carrier-mediated saturable mechanism described in Section 2.3.3.

<sup>5</sup> 11 mg choline/1,000 kcal per day, i.e. about 2.6 mg choline/MJ per day.

### 2.3.6.1.1. Choline and trimethylamine-N-oxide (TMAO)

Excretion of choline in the urine is low in relation to usual dietary intakes. De la Huerga and Popper (1951) (Section 2.3.5.2.2) determined the excretion of choline and TMA in the urine in four healthy adult subjects after single oral doses of 2–8 g of choline (as choline bicarbonate). The authors detected no or negligible choline in urine at baseline and not more than 0.3% of the administered dose thereafter. Within 24 h, two-thirds of the administered dose was excreted as TMA and TMAO, which suggests that unabsorbed choline was metabolised by the intestinal microbiota.

In pregnant and non-pregnant women (consuming either 480 or 930 mg of choline/day for 12 weeks), the (geometric) mean of the excretion of choline in the urine throughout the 12-week study was 10.7 (95% confidence interval (CI): 8.1–14.1) and 3.2 (95% CI: 2.3–4.4) mg/day, respectively ( $p \leq 0.001$ ), and did not change significantly with choline intake (Yan et al., 2012) (Sections 2.3.6.1.2, 2.4.1.2, 5.1.3 and 6.4). In lactating and non-lactating women (the latter from the study by Yan et al. (2012)), mean excretion of choline in the urine throughout the study (10–12 weeks) did not differ (Davenport et al., 2015) (Sections 2.3.3, 2.3.6.1.2, 2.3.6.3, 2.4.1.2 and 5.1.3).

### 2.3.6.1.2. Betaine and dimethylglycine (DMG)

Betaine in the urine originates either from the diet or is formed in the kidney (and liver) via CHDH and BADH from choline. In this reaction, betaine is a methyl-group donor for Hcy remethylation (Figure 2 and Sections 2.2.2.1 and 2.3.5.2.1). BHMT demonstrates saturation kinetics, its activity increases in rat liver when the diet is low in methionine but contains choline or betaine (Park and Garrow, 1999) and its activity is inhibited by DMG, which is the product of BHMT activity. Moreover, oxidative demethylation of DMG to sarcosine is the rate-limiting step in betaine metabolism. Betaine normally accumulates in the kidney medulla, where its release into the urine is controlled by intracellular tonicity.

While the betaine plasma concentration remains almost stable on a habitual diet, it increases rapidly about 30-fold following one oral dose of about 50 mg betaine/kg bw in 12 healthy males and has an elimination half-life of around 14 h (Schwahn et al., 2003a). In this study, on average, 4% of the ingested dose was excreted as betaine in the 24-h urine; the renal clearance<sup>6</sup> was in the range of 0.4–13.9 mL/h per kg bw and about 5% of the apparent total plasma clearance. Betaine is freely filtered in the kidney, but normally almost completely reabsorbed in the proximal tubule (Lever et al., 2007).

In a randomised cross-over study on eight healthy males consuming five different intervention meals, including one high-choline meal (564 mg) or a single dose of choline supplement (500 mg, as choline chloride), compared to a low-choline meal (< 1 mg choline), urinary betaine excretion was not significantly different between groups (Atkinson et al., 2008). In contrast, in this study, urinary DMG excretion peaked at 4–6 h ( $p < 0.005$  compared to control), but was still higher than baseline ( $p < 0.05$ ) 24 h after the high-choline meal.

In women consuming either 480 or 930 mg of choline/day for 12 weeks (Yan et al., 2012), the (geometric) mean of the excretion of betaine in the urine throughout the 12-week study was 12.9 (95% CI: 10.0–16.6) mg/day in pregnant women and 8.1 (95% CI: 6.1–10.8) mg/day in non-pregnant women ( $p \leq 0.05$ ) (Sections 2.3.6.1.1, 2.4.1.2, 5.1.3 and 6.4). Lactating women had a lower excretion of choline metabolites (betaine: –3 mg/day,  $p = 0.001$ ; DMG: –2.3 mg/day,  $p < 0.001$ ) in the urine throughout the study period compared to non-lactating control women (Davenport et al., 2015) (Sections 2.3.6.1.1, 2.3.6.3, 2.4.1.2 and 5.1.3).

Infants excrete high amounts of betaine in their urine, up to 1.5 mmol/mmol creatinine (1.55 g/g creatinine) during the first year of life, with a maximum at the age of 2–3 months and a decrease to 0.2 mmol/mmol creatinine at 1 year (Holmes et al., 1996). During the first 10 days of life, a urinary excretion of betaine of  $27.4 \pm 2.8 \mu\text{mol/kg bw per day}$  ( $3.2 \pm 0.3 \text{ mg/kg per day}$ ; mean  $\pm$  SEM) was reported in 27 infants. At that age, no dietary source of betaine is available (Holmes et al., 1996), besides human milk with its low concentration of betaine (360–515  $\mu\text{g/L}$ ) (Davenport et al., 2015). In the newborn period, urinary excretion of betaine may be higher than choline intake (Davies et al., 1992).

<sup>6</sup> Defined as the ratio of 24 h urinary excretion (mmol/kg bw) to the respective area under the curve (in mmol/L per hour).



### 2.3.6.1.3. Conclusion on urinary excretion

The Panel notes that choline excretion in the urine is low in relation to usual dietary intakes (and 0.3% of the administered dose of 2–8 g choline). A study showed that pregnant women have higher urinary excretion of choline and betaine than non-pregnant women. The Panel notes that excretion of betaine in urine may be of dietary origin or produced from choline. The rise in urinary DMG concentration, the second product of BHMT activity, after a choline supplement or a high-choline meal, suggests that choline-derived betaine is primarily used for Hcy remethylation in the liver (rather than fulfilling the other functions of betaine in the body).

### 2.3.6.2. Faeces

Hepatic PC, synthesised either from dietary choline via the CDP-choline pathway or via the PEMT pathway (Figure 2), is used for secretion of VLDL or formation of high-density lipoproteins (HDL), or secretion into the bile. In mice, PC secretion into the bile was equivalent to the entire hepatic PC pool, of which 95% is reabsorbed (Li and Vance, 2008). In addition, PC is secreted by the intestinal mucosa, according to data in animals and patients (Ehehalt et al., 2010).

No human data are available on faecal excretion of choline or choline compounds in relation to dietary choline intake. Depending on the composition of the gut microbiome, non-absorbed choline in the gut can be converted to TMA (Sections 2.2.2.2 and 2.3.5.2.2).

### 2.3.6.3. Breast milk

Choline is found in breast milk predominantly as PChol and GPC, together with free choline, PC, SPM. Its concentration changes during the progress of lactation, and is influenced by maternal diet (Fischer et al., 2010b; Davenport et al., 2015). Apart from choline and choline containing compounds, breast milk also contains betaine (Davenport et al., 2015).

In an RCT in 103 pregnant (then lactating) women (94 completers), Fischer et al. (2010b) (Sections 2.3.3, 2.4.1.2, 2.5.1, 5.1.3 and 5.2.5) investigated the response of maternal plasma and breast milk choline concentrations to a PC supplement (750 mg/day choline,  $n = 48$ , from the 18th gestational week to 90 days post partum), compared to placebo ( $n = 46$ ). The supplement was consumed in addition to a mean dietary choline intake of about 350 mg/day (measured by a three-day food record at 45 days post partum). Breast milk (and maternal plasma) concentrations were measured at 45 days post partum. There was a significant linear correlation between total choline intake (sum of all choline compounds from foods and supplements, range about 150 mg/day to > 750 mg/day) and breast milk concentrations of PChol, PC, free choline and betaine ( $R^2 = 0.16$  and  $p = 0.0001$ ,  $R^2 = 0.07$  and  $p = 0.02$ ,  $R^2 = 0.08$  and  $p = 0.001$ ,  $R^2 = 0.13$  and  $p = 0.0003$ , respectively), when all subjects were taken into account. Mean ( $\pm$  SE) breast milk concentrations of PChol ( $722 \pm 39$  vs  $553 \pm 27$   $\mu\text{mol/L}$ ) and free choline ( $106 \pm 10$  vs  $83 \pm 8$   $\mu\text{mol/L}$ ) were significantly higher ( $p < 0.001$ ) in the supplemented group than in the placebo group, whereas PC, GPC and SPM were not significantly different.

In a controlled feeding study, Davenport et al. (2015) (Sections 2.3.3, 2.3.6.1, 2.4.1.2 and 5.1.3) investigated the response of breast milk choline concentration to different choline intakes. In this study, lactating ( $n = 28$ , 5 weeks post partum) and non-pregnant non-lactating ( $n = 21$ , control) women were randomised to consume 480 mg/day (15 lactating women and 10 controls) or 930 mg choline/day (13 lactating women and 10 controls), from food and supplements,<sup>7</sup> for 10 (lactating women) or 12 weeks (control women). Lactating women consuming 930 mg/day choline had a significantly higher concentration of total choline in breast milk (sum of all choline compounds) at the end of the study compared to those consuming 480 mg/day (mean  $\pm$  SD:  $1,200 \pm 60$  vs  $1,000 \pm 50$   $\mu\text{mol/L}$ ,  $p = 0.041$ ). They also had higher concentrations of PChol ( $392 \pm 26$  vs  $285 \pm 24$   $\mu\text{mol/L}$ ,  $p = 0.008$ ) and GPC ( $471 \pm 36$  vs  $346 \pm 33$   $\mu\text{mol/L}$ ,  $p = 0.031$ ), but their free choline concentration in breast milk did not differ ( $148 \pm 13$  vs  $158 \pm 12$   $\mu\text{mol/L}$ ). During the last 4–6 weeks, 20% of the total intake of choline was provided as deuterium labelled choline (methyl-D<sub>3</sub>-choline). Women consuming the higher choline intake (930 mg/day) during lactation had in their breast milk, at the end of the study, a significantly higher enrichment of the metabolites generated endogenously via the hepatic PEMT pathway, but not of the metabolites generated from intact exogenous choline via the CDP-choline pathway (see pathways in Figure 2, Section 2.3.5). The

<sup>7</sup> Diet provided an average of 380 mg/day of choline, and supplemental choline was 100 or 550 mg/day.

Panel notes that the higher choline intake during lactation (930 mg/day compared to 480 mg/day) significantly increased the concentration of total choline in breast milk, and increased the supply of PEMT-derived choline metabolites in breast milk.

The content of PC and SPM in breast milk was reported to remain constant from day zero to 85 of lactation, while the content of GPC, PChol and, to a lesser extent, free choline, in breast milk increased significantly after the first week after birth (Zeisel et al., 1986), but only free choline content decreased significantly with time.

A search of the literature published after January 2000 was performed as preparatory work to this assessment, in order to identify breast milk composition data for choline (LASER Analytica, 2014). This search was completed with two additional papers (Holmes-McNary et al., 1996; Davenport et al., 2015). Appendix A reports data from six studies (Holmes-McNary et al., 1996; Holmes et al., 2000; Ilcol et al., 2005; Fischer et al., 2010b; Ozarda et al., 2014; Davenport et al., 2015) conducted in the UK, Turkey and the USA, on the mean/median free and total choline concentrations of human milk from healthy lactating mothers. Either the infants were full-term (Holmes-McNary et al., 1996; Ozarda et al., 2014), or there was a mixed population of full-term and preterm infants or it was unclear whether the infants were born at term or not.

Stages of lactation varied between birth and 180 days post partum. Mean maternal choline intake was not reported in four studies (Holmes-McNary et al., 1996; Holmes et al., 2000; Ilcol et al., 2005; Ozarda et al., 2014), while one study compared choline supplemented versus non-supplemented women (Fischer et al., 2010b) and the other compared two doses of choline supplementation (Davenport et al., 2015). Three studies (Ilcol et al., 2005; Fischer et al., 2010b; Davenport et al., 2015) reported information on maternal plasma choline concentration (considered by the authors as an indication of maternal status). The mean/median concentration of total choline (sum of compounds) in mature milk ranged from 120 to 160 mg/L (see Appendix A).

Based on the two studies on full-term fully breast-fed infants (Holmes-McNary et al., 1996; Ozarda et al., 2014) in the USA and Turkey (n = 70 women in total), an average total choline concentration (free choline and choline compounds) of about 145 mg/L in mature breast milk can be calculated. Assuming a mean milk transfer of 0.8 L/day during the first 6 months of lactation in exclusively breastfeeding women (Butte et al., 2002; FAO/WHO/UNU, 2004; EFSA NDA Panel, 2009), the estimated secretion of choline into milk during lactation would be 116 mg/day, rounded to 120 mg/day.

The Panel notes that breast milk mainly contains PChol and GPC, besides free choline, PC and SPM, in concentrations depending on the progress of lactation and maternal diet/supplementation. The Panel also notes that increased maternal choline intake enhances the concentration of total choline in breast milk and increases the supply of PEMT-derived choline metabolites in breast milk. The Panel considers that secretion of choline into breast milk during the first 6 months of exclusive breastfeeding is about 120 mg/day.

### 2.3.7. Interaction with other nutrients: folate

Folate and choline are both involved in the remethylation of Hcy to methionine, the first via 5-methyl-THF, the latter via betaine. The interrelationship between folate and choline metabolism has been demonstrated in animal studies (Varela-Moreiras et al., 1992; Kim et al., 1994) (Section 2.3.5.2.1). In the first case, Hcy is methylated to methionine by the ubiquitous methionine synthase (MS, Figure 2), which requires methyl-THF as methyl-group donor and cobalamin as cofactor (Ueland et al., 2005). In the second case, Hcy is methylated to methionine by BHMT (Figure 2), which requires betaine as methyl-group donor. Choline insufficiency, with consequently low betaine formation, increases the requirement for methyl-THF for the remethylation of Hcy and, therefore, the requirement for dietary folate. Vice versa, in folate depletion, methyl groups from choline and betaine are increasingly used for Hcy remethylation, thereby increasing the requirement for choline. Methyl-THF and choline/betaine can be considered as partially exchangeable sources of methyl groups (Kim et al., 1994).

Jacob et al. (1999) investigated the effect of folate depletion and repletion on choline status and the *in vivo* methylation capacity in humans residing in metabolic units. Following a baseline period of 6–9 days on a diet sufficient in energy and all nutrients including folate (440 µg/day), 11 healthy men (aged 33–46 years) and 10 healthy women (aged 49–63 years) consumed, for 4–5 weeks, a low folate (average of 25 µg/day and 56 µg/day for men and women, respectively) and low choline diet (average of 238 mg/day and 147 mg/day for men and women, respectively). Two to six weeks of folate repletion followed (440 and 516 µg folate/day for men and women, respectively, partially supplied as folic acid) without change in the choline intake. Variation in the methionine content of the diet in men

(400 mg or 1,400 mg/day in the first half of the study period with cross-over thereafter) had no effect on the outcomes (this was not investigated in women). No functional deficiencies of organs were noted in any subject. Methylation capacity, as assessed by the urinary excretion of creatinine and of methylated nicotinamide breakdown products after ingestion of 1 g of nicotinamide, was not diminished. At the end of the folate depletion phase, plasma free choline (and folate) concentrations were significantly lower in both men and women compared with baseline, and plasma tHcy concentration was significantly higher while plasma PC concentration was decreased in men compared to baseline (PC concentration was not investigated in women). At the end of the folate repletion phase, plasma choline concentrations increased significantly in both sexes compared to the folate depletion phase ( $p < 0.05$ ), in women to even higher values than at baseline ( $p < 0.05$ ), with no significant change in plasma tHcy concentration compared to the folate depletion phase. No changes in choline, folate and SAM concentrations in red blood cells were noted throughout the study. The Panel notes that, in this study, an adequate folate intake maintained plasma choline concentration despite a low choline intake of about 150–250 mg/day on average, while plasma free choline and PC concentrations decreased and tHcy concentration increased when both folate and choline intakes were low.

In 43 premenopausal Mexican-American women, folate intake was restricted for 7 weeks to 135 µg dietary folate equivalent (DFE) per day, followed by 7 weeks of randomisation to either 400 or 800 µg DFE/day, while choline intake was kept constant at 349 mg/day (including 250 mg/day of a choline supplement) (Abratte et al., 2008). In this study, plasma PC concentration decreased during dietary folate restriction compared with baseline ( $p = 0.001$ ), presumably due to the unfulfilled demand of folate-derived one-carbon units for PC synthesis. Plasma PC concentration increased again after administration of 800 µg DFE/day ( $p = 0.03$ ) (but not significantly with 400 µg DFE/day). The Panel notes that, in this study, folate intake was shown to influence plasma PC concentration.

Changes in the activity of enzymes involved in folate and choline metabolism, due to polymorphisms of genes for enzymes of this metabolism, can be expected to have an impact on the status of folate and choline. An example is the C677T genotype of the MTHFR (Sections 2.3.5.2.1 and 2.5), which has a strong influence on folate status (Abratte et al., 2008).

Ivanov et al. (2009) examined the potential influence of polymorphisms of two genes involved in choline metabolism (*MTHFD1* rs2236225 and *PEMT* rs12325817 and rs7946) (Section 2.5. and Appendix C) on plasma PC and tHcy concentrations in the presence of folate restriction in the same Mexican-American women studied by Abratte et al. (2008). These polymorphisms are functional in that they impair the activity of the two enzymes (*PEMT* and *MTHFD1*) and thereby possibly increase choline requirement and compromise the production of methyl-THF. The *PEMT* and *MTHFD1* polymorphisms did not modify the small negative response of plasma PC concentration to folate restriction, except in case of homozygosity for *PEMT* rs1232587 that attenuated the decline in plasma PC concentration. Homozygosity for *PEMT* rs7946 and *MTHFD1* rs2236225 SNPs was associated with a greater increase ( $p < 0.001$ ) in plasma tHcy concentration during folate restriction than in subjects homozygous for the wild-type.

The Panel notes that, in the available studies, low folate intake had a negative impact on plasma PC concentration in the presence of 'adequate' choline intake, and that the impact of SNPs of genes of some enzymes involved in metabolic pathways of choline may result in increased tHcy concentrations in plasma during folate restriction. These changes are not predictable due to compensatory changes in other parts of those pathways. The Panel, moreover, notes the small number of subjects investigated and stratified for genetic polymorphisms that limits the generalisation of these studies.

## 2.4. Biomarkers

### 2.4.1. Plasma/serum concentration of free choline and choline-compounds and metabolites

#### 2.4.1.1. Adults

Fasting plasma free choline concentrations usually range between 7 and 20 µmol/L, with most subjects having a concentration of 10 µmol/L (IOM, 1998). Plasma choline concentrations are regulated, however, some variability in plasma concentrations occurs with changes in choline intake. Choline-deficient diets, as applied in depletion/repletion studies (Section 5.1.2) and consumed over weeks, can reduce plasma free choline concentration by approximately 30%, and ingestion of choline-rich foods (e.g.  $\geq 500$  mg/day) can increase plasma free choline concentrations beyond 20 µmol/L (Zeisel et al., 1991) (Appendix D). Plasma free choline concentration was found not to decrease

beyond 50% of the initial normal value even after 1 week of total fasting, presumably because of release of choline from membrane phospholipids (Savendahl et al., 1997).

Fasting plasma PC concentration varied between adults (1.5–2.5 mmol/L) and decreased by 30% after 3 weeks on a low choline diet, while erythrocyte PC concentration decreased by 10% (Zeisel et al., 1991).

#### 2.4.1.2. Pregnancy and lactation

During pregnancy, serum free and phospholipid-bound choline concentrations increase compared with non-pregnant women (Ozarda Ilcol et al., 2002).

The controlled feeding study by Yan et al. (2012) (Sections 2.3.6.1, 5.1.3 and 6.4) compared the effects of two doses of choline supplementation (480 or 930 mg of choline/day from food and supplements) in healthy pregnant (recruited at 27 weeks gestation) and non-pregnant women. In this study, pregnant women had similar mean plasma free choline concentration as non-pregnant women at recruitment, but significantly higher concentration (by 30%) than non-pregnant women throughout the 12-week study (geometric means, (95% CI): 8.2 (7.6–8.7) vs 6.3 (5.6–6.9)  $\mu\text{mol/L}$ , respectively,  $p < 0.001$ ). Pregnant women had lower mean plasma concentrations of the three methyl-group donors (betaine, DMG, sarcosine) as well as methionine and Hcy at recruitment, and this persisted throughout the study (lower by 13–55%,  $p < 0.001$ ). The lower circulating concentrations of choline-derived methyl-group donors in pregnant women, than in non-pregnant women, throughout the study, were possibly a consequence of the greater use of these molecules in both maternal and fetal compartments. Pregnant women consuming 930 mg choline/day had a 13% higher mean plasma concentration of free choline than those consuming 480 mg choline/day ( $p = 0.021$ ).

In a prospective observational study, choline intake of 154 pregnant women, estimated by a food frequency questionnaire (FFQ), was weakly correlated to their natural log-transformed plasma concentration of free choline at 16 and 36 weeks of gestation (16 weeks:  $r = 0.20$ ,  $p = 0.013$ , range of intake read on figure: 150–700 mg/day) (Wu et al., 2012).

In a prospective cohort study on 368 Canadian pregnant women recruited at 12–16 weeks of gestation, Visentin et al. (2015) investigated the relationship between maternal choline intake and concentrations of choline and its metabolites in maternal and umbilical cord plasma. Mean maternal choline intake (total of all compounds), as estimated by a semiquantitative FFQ, was  $306 \pm 127$  and  $302 \pm 122$  mg/day in early (0–16 weeks) and late (23–37 weeks) pregnancy, respectively. Mean maternal plasma free choline (95% CI) was 7.2 (7.1–7.4)  $\mu\text{mol/L}$ . The mean concentrations of free choline, DMG and TMAO in maternal plasma increased significantly ( $p \leq 0.005$ ) between recruitment in pregnancy and delivery by 49%, 17% and 13%, respectively, whereas that of betaine decreased by 21% ( $p \leq 0.005$ ). Maternal dietary intake (total or free) was not associated with these maternal plasma concentrations. The mean concentrations of free choline, betaine and DMG in cord plasma were 3.2, 2.0 and 1.3 times the concentrations in maternal plasma at delivery, whereas the mean concentration of TMAO in cord plasma was lower by 12%. Maternal dietary choline intake (or fetal genetic variants in genes involved in choline metabolism,<sup>8</sup> Appendix C) was not associated with cord plasma concentrations of free choline and its metabolites. In contrast, maternal plasma concentrations of betaine, DMG and TMAO at delivery strongly influenced umbilical cord plasma concentrations ( $r^2$  between 0.19 and 0.51, all  $p < 0.0001$ , after adjustment for potential confounders). There was only a weak correlation between the concentration of free choline in maternal and umbilical cord plasma ( $r^2 = 0.12$ ,  $p = 0.06$ ).

Results are indicative of an active transport of choline from the mother to placental tissue (Section 2.3.3) and/or an uptake and metabolism of choline by the fetus reflecting a demand of the fetus for choline and methyl-group donors.

In lactating women, serum free and phospholipid-bound choline concentrations were significantly higher than in non-lactating women ( $p < 0.05$ ), and gradually decrease until 180 days after the birth of the child (Ilcol et al., 2005).

In the lactating women of the RCT by Fischer et al. (2010b) (Sections 2.3.3, 2.3.6.3, 2.5.1, 5.1.3 and 5.2.5), there was a significant correlation between total choline intake (from foods and supplements) and maternal plasma concentration of free choline ( $R^2$  of 0.15 in the supplemented group, and 0.55 in all subjects combined,  $p = 0.03$  and  $p = 0.0001$ , respectively). Choline supplementation increased mean maternal plasma concentration of free choline compared to placebo (mean  $\pm$  SE:  $13.7 \pm 0.6$  vs  $7.7 \pm 0.3$  nmol/mL at 45 days post partum,  $p < 0.001$ ).

<sup>8</sup> Ten SNPs in seven candidate genes.



In addition, in the controlled feeding study by Davenport et al. (2015) (Sections 2.3.3, 2.3.6 and 5.1.3), lactating women showed higher (+ 27%,  $p < 0.001$ ) plasma free choline concentrations than non-pregnant non-lactating women throughout the study period. Lactating women who consumed 930 mg/day choline had significantly higher plasma free choline concentration (+ 16%,  $p = 0.012$ ) compared to those consuming 480 mg/day.

#### 2.4.1.3. Infants

In newborns, serum free choline concentrations were significantly higher (greater than twice maternal values) and phospholipid-bound choline concentrations were significantly lower (by about 40%) than in their mothers (Holmes et al., 2000). Phospholipid-bound choline plasma concentrations in the infants rose by 40% starting from day 5 to day 15 after birth to reach adult levels by the age of about 10 years. Plasma free choline concentration of newborns remained high for 2 weeks after birth, was still slightly higher than adult levels at the age of 2 years and remained stable at around 10  $\mu\text{mol/L}$  at the age 3–12 years. This high newborn's plasma concentration possibly reflects the increase in choline in breast milk in the second week of life (Section 2.3.6.3). There was no correlation between maternal and newborn plasma phospholipid-bound choline (Buchman et al., 2001; Icol et al., 2005).

#### 2.4.1.4. Conclusion on plasma/serum concentration of choline and choline-compounds and metabolites

The Panel notes age-related changes in choline concentrations in plasma/serum, with higher values in infants and young children than in adults. Plasma choline concentration may increase when intake is increased, and decreases by up to 50% when dietary intake is severely restricted (Zeisel et al., 1991; Savendahl et al., 1997).

The Panel also notes that pregnancy and lactation are associated with higher free choline concentrations in plasma than in the non-pregnant non-lactating state, and that choline supplementation increases maternal plasma concentration of free choline in pregnancy or lactation. However, the Panel considers that the maternal intake of choline cannot be deduced from the choline concentration in maternal plasma during early and late pregnancy or lactation, nor from the choline concentration in venous umbilical cord plasma.

No dose–response relationship between choline intake and plasma concentration of free choline, or of PC, betaine, DMG or TMAO, or erythrocyte PC, can be deduced from the available data. Therefore, the Panel considers that plasma concentrations of choline and choline compounds or metabolites cannot be used for setting DRVs for dietary choline.

#### 2.4.2. Total trimethylamine (TTMA) hepatic production

The Panel concludes that TTMA hepatic production and excretion in urine are not predictably related to dietary choline intake and cannot be used for setting DRVs for dietary choline (Section 2.3.5.2.2).

#### 2.4.3. Plasma total homocysteine

Appendix B compiles the results of six studies on adults (19–82 years of age) investigating the influence of choline intake on plasma tHcy concentrations. Three studies were RCTs (Olthof et al., 2005; Atkinson et al., 2008; Wallace et al., 2012), with choline given as supplements (500–2,600 mg/day of choline) for 2–12 weeks or just once a week. Three others were cross-sectional studies within long-term cohorts (Cho et al., 2006; Chiuve et al., 2007; Lee et al., 2010b), involving 6,069 subjects of which 1,325 were men. The results from RCTs with supplements are inconsistent. RCTs with choline doses of 500 and 1,000 mg/day showed no decrease in plasma tHcy concentration (Atkinson et al., 2008; Wallace et al., 2012). However, a dose of 2,600 mg/day (as PC) over 2 weeks resulted in a significant decrease in fasting plasma tHcy concentration (mean  $\pm$  SD:  $15.6 \pm 4.0$  vs  $13.6 \pm 2.5$   $\mu\text{mol/L}$ ;  $p < 0.0001$ ) and, compared to placebo, a significantly lower rise ( $p < 0.0001$ ) in plasma tHcy concentration following a methionine load (0.1 g/kg bw) (Olthof et al., 2005). The cross-sectional studies showed an inverse relationship between dietary choline intake (that ranged in quintiles from around 230 to 400 mg/day) and fasting plasma of tHcy concentrations.

The Panel notes that many factors besides dietary or endogenous choline determine the tHcy concentration in plasma. The Panel concludes that neither fasting nor post-methionine load tHcy concentrations in plasma can be used for setting DRVs for dietary choline.

#### 2.4.4. Urinary betaine excretion

The Panel notes that betaine in urine may be of dietary origin or produced in the body from choline (Section 2.3.6.1.2). The rise in urinary DMG concentration, the second product of BHMT activity (Figure 2 and Section 2.3.6.1.3), after a choline supplement or a high-choline meal, suggests that choline-derived betaine is primarily used for Hcy remethylation in the liver (rather than fulfilling the other functions of betaine in the body).

The Panel concludes that urinary betaine excretion is not predictably related to dietary choline intake and, therefore, cannot be used for setting DRVs for dietary choline.

#### 2.4.5. Conclusions on biomarkers

The Panel considers that the available data do not allow concluding on a dose–response relationship between choline intake and plasma choline concentration or the use of this concentration as a biomarker of choline status. The Panel also considers that plasma free choline concentrations are not suitable to derive DRVs for dietary choline. Plasma concentrations of PC, betaine, DMG, tHcy or TMAO, erythrocyte PC concentration, or urinary betaine and TTMA excretion can neither be used to set DRVs for dietary choline.

### 2.5. Effects of genotypes involved in choline metabolism

Several SNPs in genes coding for enzymes in choline metabolism and in methyl-group metabolism can alter the requirement for choline and determine the likelihood of developing signs of choline deficiency (Section 2.2.2.1) with low dietary choline intakes. For example, MTHFD1 (Sections 2.2.2.1 and 2.3.7) is a trifunctional enzyme responsible for generating and interconverting 1-carbon-substituted THF cofactors from formate. *MTHFD1* mutations can impact both Hcy remethylation and thymidylate (dTMP) biosynthesis.

Genetically modified mice with defective MTHFR activity become choline deficient (Schwahn et al., 2003b) and 15–30% of humans have genetic polymorphisms that alter the activity of this enzyme resulting in a higher requirement for folate, and potentially indirectly for choline if folate intake is lower than the requirement (Rozen, 1996; Wilcken et al., 1996).

da Costa et al. (2006a) (Sections 2.2.2.1. and 5.1.1.3 and Appendices C and D) performed a controlled trial in 57 subjects (26 men and 31 women), aged 18–70 years, to determine whether susceptibility to develop organ dysfunction due to choline deficiency was influenced by common genetic polymorphisms. The choline depletion/repletion study design is described in Section 5.1.1.2. Sixty-eight percent of the subjects ( $n = 39$ ) developed organ dysfunction on the low-choline diet, which was resolved during choline repletion. Mean plasma choline concentrations decreased by almost 30% (from 9.8 to 7.1  $\mu\text{mol/L}$ ), irrespective of development of organ dysfunction. Susceptibility to choline deficiency was not affected by *BHMT* +742G→A SNP (rs3733890) in this study.

Niculescu et al. (2007) (Section 5.1.1 and Appendix D) performed a study in 33 subjects (14 men and 19 women), aged 20–67 years, to examine the effects of a low-choline diet on gene expression in subjects who developed organ dysfunction due to low choline intake, those who did not, and the potential role of four SNPs in genes involved in folate and choline metabolism (*PEMT* rs12325817, *MTHFD1* rs2236225, *CHDH* rs9001 and rs12676). The choline depletion/repletion study design is described in Section 5.1.1.2. Blood was collected after the baseline diet and after the low-choline diet, and peripheral lymphocytes were used to measure gene expression and for SNP genotyping. The low-choline diet resulted in underexpression of 152 genes and overexpression of 107 genes. Differences in gene expression changes were noted between those who developed organ dysfunction and those who did not. Analyses using group clustering and gene ontology showed that changes in gene expression related to the experimental diets were significantly altered by the SNPs examined.

Appendix C lists the enzymes (*PEMT*, *MTHFD1*, *CHDH*, *BHMT*, choline kinase isoform A or B (*CHKA* or *CHKB*), *CCT*, *SLC44A1*, *MTHFR*), which have SNPs with known qualitative impact on choline requirement and/or are associated with an increased risk of developing organ dysfunction or other health outcomes, including birth defects, when consuming a low-choline diet. In particular, some specific polymorphisms of the genes for *PEMT*, *CHDH* and *MTHFD1* were shown to increase the dependency on dietary choline intake (Appendix C).

According to the review by Au et al. (2010), it may not be accurate to include or exclude risk contribution of the tested genes investigated in epidemiological studies on neural tube defects (NTDs), some of them having limitations in study design, that potentially affect the power of statistical analysis,

thus providing conflicting conclusions. For complex diseases like NTDs, it is anticipated that the risk of a disease-associated allele is between 1 and 2, and over 2,000 samples (cases plus controls) would be needed to provide statistical power of 80% to assess a risk of 1.8–2 of a disease locus with a SNP allele frequency of 0.1. Double or quadruple the controls would be needed if unmatched controls are used, to adjust for confounding factors.

The Panel notes that many SNPs have been described for genes coding for eight enzymes (Appendix C) involved in choline or methyl-group metabolism, and that carrier frequency in mixed populations can be up to about 70%. Kohlmeier et al. (2005) mention a personal communication by K. Meyer and P.M. Ueland that the distribution of polymorphic variants of *MTHFR* and *MTHFD1* in North Carolina (Appendix C) largely agreed with that of North European populations (Norwegian Colorectal Cancer Prevention (NORCCAP) study).

The effects of the *PEMT* polymorphism rs12325817, on the likelihood of development of signs of organ dysfunction (mainly liver) when choline intake is experimentally restricted to  $\leq 50$  mg/70-kg bw per day, have been investigated most often. The risk of organ dysfunction is higher in postmenopausal than in premenopausal women and is increased by simultaneous restriction in folate intake. Due to the experimental design of choline depletion/repletion studies with a low choline intake during the depletion period ( $\leq 50$  mg/70-kg bw per day) (Section 5.1.1), and because of the lack of data on the relationship between habitual choline intakes and signs attributable to choline deficiency in populations, the Panel notes that the amount of dietary choline needed to prevent such signs cannot be predicted with confidence.

### 2.5.1. Influence of polymorphisms in pregnancy and lactation

Polymorphisms in the *MTHFD1* gene and the *BHMT* gene, coding for enzymes involved in choline metabolism, were identified as potential candidates for association with choline concentrations in maternal plasma and breast milk (Fischer et al., 2010b) (Sections 2.3.3, 2.3.6.3, 2.4.1.2, 5.1.3, 5.2.5).

In the RCT by Fischer et al. (2010b), the authors also investigated whether maternal polymorphisms (370 SNPs in 10 genes involved in choline metabolism) modified the response of maternal plasma and breast milk choline concentrations (measured at 45 days post partum) to choline supplementation (compared with placebo). These SNPs were tested in linear regression models, with choline metabolites as the response and homozygous wild-type, heterozygous wild-type and homozygous variant alleles of SNPs, as well as choline intake (from food and supplements), as predictors. In these models, five SNPs in the *MTHFR* gene were identified in the placebo group that, for most of them, reduced the slope of the response curve of free choline concentration in breast milk to choline intake ( $p < 0.05$ ). In addition, outliers previously identified by the authors (in a first analysis of the relationship between intake and concentrations in breast milk or plasma) were tested for combinations of shared SNPs. In this analysis, three subjects of the placebo group were identified with five SNPs in common in the *MTHFD1* gene and who had exceptionally high breast milk choline concentrations (in relation to choline intake). Five participants were also identified with two SNPs in common in the *BHMT* gene, and four of these subjects had lower than average plasma free choline concentrations (in relation to choline intake).

Besides the choline intake of the mother (Section 2.3.6.3 and Appendix A), the Panel notes that polymorphisms in genes coding for enzymes involved in choline and methyl-group metabolism, particularly if they occur in combinations, can influence the amount of choline secreted into breast milk. The Panel considers that the available data on polymorphisms in genes are insufficient to predict choline concentrations in breast milk.

### 2.5.2. Conclusion on effects of genotypes

The Panel concludes that SNPs can enhance or reduce the function of enzymes involved in choline metabolism. This can influence the requirement for choline and, moreover, can determine the susceptibility to dietary choline deficiency. The Panel considers that particularly some specific polymorphisms of the genes for the enzymes *PEMT*, *CHDH* and *MTHFD1* are known to increase the dependency on dietary choline intake. Since their frequency in populations vary and their impact on dietary choline requirement may be influenced by dietary habits, no conclusions can be drawn from available studies on predictable variations in individual choline requirements.



### 3. Dietary sources and intake data

#### 3.1. Dietary sources

Total choline content is highest in eggs (raw egg yolk: about 670 mg/100 g food, whole raw fresh egg: about 290 mg/100 g food) followed by meats and fish, whole grains, vegetables and fruit, and fats and oils (median content of fats and oils: about 5 mg/100 g food) (USDA, 2015) (Section 3.2.1). The proportion of different choline compounds in food can change by preparation. For example, cooking decreases the concentration of free choline and increases the content of PC per 100 g food, while mincing of raw vegetables decreases the content of PC by activating phospholipase D with the release of free choline and phosphatidic acid (Zeisel et al., 2003). The implications of such changes in choline compounds for human nutrition are unknown.

Human milk is rich in choline (Section 2.3.6). Ilcol et al. (2005) showed that the distribution of choline compounds in human milk, and bovine-derived and soy-protein based formulae from different manufacturers differed considerably, e.g. soy-derived formulas had much less SPM than human milk.

In the EU, the addition of choline to infant formula is mandatory with a minimum level of 7 mg and a maximum level of 50 mg of choline/100 kcal and the total phospholipid concentration must be not higher than 2 g/L.<sup>3</sup>

Currently, choline, choline chloride, choline bitartrate and choline citrate may be added to food intended for infants and young children, food for special medical purposes, and total diet replacement for weight control in the EU.<sup>9</sup> CDP-choline (citicoline) has been evaluated as novel food by EFSA and no safety concerns were raised (EFSA NDA Panel, 2013a).<sup>10</sup> Choline and choline compounds can be found in dietary supplements.

#### 3.2. Dietary intake

##### 3.2.1. Dietary intake in EU countries

The Panel notes that no food composition data with respect to choline are available at the European level, and that there is a lack of reliable measurements of choline content in foods in the EU. The Panel refers to the study by Vennemann et al. (2015), which used, with the aim at assessing choline intake in the EU, the total choline composition data from the release 26 of the National Nutrient Database for Standard Reference from the US Department of Agriculture (USDA database; issued in November 2013) (USDA, 2013) (Section 3.1). Total choline content of US foodstuffs was calculated by USDA as the sum of free choline, GPC, PChol, PC and SPM.

In the assessment by Vennemann et al. (2015), food consumption data from the EFSA Comprehensive European Food Consumption Database (EFSA, 2011), classified according to FoodEx2 classification, were used. This assessment includes food consumption data from 12 dietary surveys from nine EU countries (Finland, France, Germany, Ireland, Italy, Latvia, the Netherlands, Sweden and the United Kingdom). These surveys used 3–7-day food records, 24-h recalls performed on at least 2 days or 48-h recalls. Individual data from these nationally representative (except for the Finnish surveys in children) surveys undertaken between 2000 and 2011 were available to EFSA. In this assessment by Vennemann et al. (2015), the nutrient composition data was obtained for 2,684 food items by recoding the USDA nutrient composition food list (based on the LanguaL food description thesaurus) to FoodEx2 classification (used for the food consumption data). Nutrient intake calculations were performed only on subjects with at least two reporting days. Choline intake from dietary supplements was not assessed. Mean, medians, 5th and 95th percentiles of intake of the population, per survey, age, class and sex, were calculated.

Data were available from four surveys for children aged from 1 to < 3 years, from seven surveys for older children, and from eight surveys for adults (including one survey during pregnancy). Total choline intake mean estimates ranged from 151 to 210 mg/day in children aged from 1 to < 3 years, 177–304 mg/day in children aged from 3 to < 10 years, 244–373 mg/day in children aged from 10 to

<sup>9</sup> Regulation (EU) No 609/2013 of the European Parliament and of the Council of 12 June 2013 on food intended for infants and young children, food for special medical purposes, and total diet replacement for weight control and repealing Council Directive 92/52/EEC, Commission Directives 96/8/EC, 1999/21/EC, 2006/125/EC and 2006/141/EC, Directive 2009/39/EC of the European Parliament and of the Council and Commission Regulations (EC) No 41/2009 and (EC) No 953/2009, OJ L 181, 29.6.2013, p. 35.

<sup>10</sup> Commission Implementing Decision 2014/423/EU of 1 July 2014 authorising the placing on the market of citicoline as a novel food ingredient under Regulation (EC) No 258/97 of the European Parliament and of the Council, OJ L 196, 3.7.2014, p. 24.

< 18 years. Total choline intake mean estimates ranged from 269 to 468 mg/day in adults aged from 18 to  $\geq$  75 years, i.e. from 332 to 468 mg/day in men and from 269 to 404 mg/day in women of this age range, respectively. From one survey in Latvia, the total choline intake mean estimate was 336 mg/day in pregnant adolescents and 356 mg/day in pregnant women.

Data on infants (< 1-year-old) were available from three out of the seven surveys, namely from Finland, Germany and Italy<sup>11</sup> (data not shown in the study by Vennemann et al. (2015)). The total choline intake mean estimates in infants ranged from 75 to 127 mg/day. The Panel notes the limitations in the methods used for assessing breast milk consumption in infants and related uncertainties in the choline estimates for infants.

Choline intake estimates are also available from a convenience sample of Flemish women (aged 18–35 years) (Pauwels et al., 2015). In this study, food consumption was assessed by FFQs covering 51 food items that had been selected because they were part of the Belgian diet and/or were the main contributors for one of four methyl-group donors (including choline), and the USDA database was also used as food composition database for choline. Despite important methodological differences with the intake assessment described above from the study by Vennemann et al. (2015), and the specific population group investigated, choline intake estimates in Flemish women (mean  $\pm$  SD: 286.6  $\pm$  105.1 mg/day) were in the same order of magnitude of the estimates produced by Vennemann et al. (2015) for several EU countries.

### 3.2.2. Dietary intake in non-EU countries

In view of the limited data on choline intake published in the EU, the Panel again refers to the study by Vennemann et al. (2015), which compared their estimates with four studies carried out in non-EU countries in adult men and women in the USA, New Zealand and Taiwan (Chu et al., 2012; USDA, 2012; Mygind et al., 2013), and pregnant and lactating women in Canada, followed from the first or second trimester to 3 months post partum (Lewis et al., 2014). Two of these studies used nationally representative data (Chu et al., 2012; USDA, 2012), all studies used 24-h recalls or three-day food records as dietary assessment methods (but not FFQs), were cross-sectional (apart from the study on pregnant and lactating women) and used the same composition database (USDA database) as Vennemann et al. (2015) although from different releases.

The mean choline intake estimates in adults was 415 and 279 mg/day in US men and women, respectively (USDA, 2012), 316 mg/day in women aged 18–40 years in New Zealand (Mygind et al., 2013) and 372 and 265 mg/day in men and women aged 18–64 years, respectively, in Taiwan (Chu et al., 2012). The mean ( $\pm$  SD) choline intake in pregnant and lactating women in Canada ranged between 340  $\pm$  148 in the second trimester and 346  $\pm$  151 mg/day at 3 months post partum (Lewis et al., 2014).

### 3.2.3. Conclusion on dietary intake

The Panel notes that mean choline intake estimates in adults ranged from 269 to 468 mg/day in national surveys from seven EU countries (Vennemann et al., 2015), was about 290 mg/day in one EU country (Pauwels et al., 2015), and were between 265 and 415 mg/day in three studies conducted in non-EU countries (Chu et al., 2012; USDA, 2012; Mygind et al., 2013). The Panel also notes that mean choline intake was about 350 mg/day in the only EU survey on pregnant women considered in Vennemann et al. (2015), as well as in one study on pregnant or lactating women in one non-EU country (Lewis et al., 2014). The Panel concludes that the choline intake data resulting from the assessment by Vennemann et al. (2015) in EU countries are generally of the same magnitude as the intakes of the published studies available in adults in EU (Pauwels et al., 2015) and non-EU countries (Chu et al., 2012; USDA, 2012; Mygind et al., 2013; Lewis et al., 2014).

## 4. Overview of dietary reference values and recommendations

To date, DRVs for choline have only been proposed by IOM (1998).

<sup>11</sup> The proportions of breast-fed infants were 58% in the Finnish survey, 40% in the German survey, 44% in the Italian survey. Most infants were partially breast-fed. For the Italian and German surveys, breast milk intake estimates were derived from the number of breastfeeding events recorded per day multiplied by standard breast milk amounts consumed on an eating occasion at different ages. As no information on the breastfeeding events was reported in the Finnish survey, breast milk intake was not taken into consideration in the intake estimates of Finnish infants.

## 4.1. Adults

IOM (1998) set Adequate Intakes (AIs), since data were not sufficient for deriving an Estimated Average Requirement (EAR) and a Recommended Dietary Allowance (RDA). The AIs for choline were based on data on the prevention of liver damage, as assessed by measuring serum ALT concentrations. The estimate was considered by the IOM as being uncertain because it was based on a single RCT by Zeisel et al. (1991) (depletion/repletion study, Section 5.1 and Appendix D). This study examined serum ALT activity in 16 healthy male hospitalised volunteers. They were supplemented with 500 mg choline/day for 1 week, then randomised to receive for three additional weeks either the choline-supplemented diet (control group,  $n = 7$ ) or the same diet without choline but with cellulose as placebo ( $n = 8$ ), then all subjects consumed the choline-supplemented diet during the fifth week of the study. A choline intake of 500 mg/day, which is approximately 7 mg/kg bw per day using the mean body weight for the control group, i.e. 74.4 kg, prevented ALT abnormalities in these healthy men. Thus, the AI was set at 550 mg/day after rounding, considering the US reference weight of 76 kg for men (National health and nutrition examination survey (NHANES) III, 1988–1994).

The IOM noted that, at that time, no studies undertaken in healthy women following a choline deficient diet were available. However, from an intervention study (Buchman et al., 1995) on one man and three women with hepatic steatosis receiving total parenteral nutrition containing 1–4 g/day of choline chloride for 6 weeks, the IOM concluded that women were just as likely as men to develop low plasma choline concentrations and fatty liver. To set an AI for women, the IOM assumed that the data used to set an AI for men could be used, even though women may use choline more efficiently, thus the derived AI for women was set at 425 mg/day based on the US reference weight of 61 kg for women (NHANES III, 1988–1994). The IOM noted some evidence that transport across the blood–brain barrier is diminished in older adults (60–85 years, compared to younger adults aged 20–40 years), suggesting the possibility of a higher requirement than for younger adults (Cohen et al., 1995). Nevertheless, for older adults, no adjustment was made to the AI.

## 4.2. Infants and children

For breast-fed infants from birth to 6 months, IOM (1998) set an AI of 125 mg/day. This AI was based on an average breast milk consumption of 0.78 L/day (Hofvander et al., 1982; Butte et al., 1984; Chandra, 1984; Neville et al., 1988; Allen et al., 1991) and an average choline concentration of 160 mg/L. This average choline concentration was obtained from 15 healthy US mothers exclusively breastfeeding and followed from 30 days up to 85 days post partum (Zeisel et al., 1986) and 33 healthy US mothers participating in the study during postnatal days 27–32 (Holmes et al., 1996). For older infants aged 7–12 months, the AI was extrapolated upwards from the AI for infants from birth to 6 months by allometric scaling and using US reference weights (NHANES III, 1988–1994), and was set at 150 mg/day. This value was confirmed by the downwards extrapolation from the AI for adults by allometric scaling using a growth factor, which gave the same result.

In the absence of data on which to base an EAR or AI for choline for children, IOM (1998) extrapolated the AIs for children aged 1–18 years from adult values, by allometric scaling using growth factors.

## 4.3. Pregnancy

IOM (1998) concluded that an increase in the AI to support pregnancy should be based on the fetal and placental accumulation of choline. The IOM took into account animal data on choline concentration in adult tissues (Pomfret et al., 1989), organ weight in the human fetus (Widdowson, 1963) and human data ( $n = 7$ ) on choline concentration in placental tissue (Welsch, 1976), and considered an average choline concentration of 321 mg/kg of fetal and placental tissue combined. The IOM assumed that there is no extra choline synthesis by the mother during pregnancy, and that there is no choline synthesis by the placenta or fetus. Thus, the required additional dietary intake of choline for 10 kg of tissue, that comprises the fetus (3 kg) and organs of pregnancy (7 kg), was calculated to be approximately 11 mg/day throughout pregnancy. The AI for choline was thus set at 450 mg/day (after rounding) for pregnant adolescent and adult women.

## 4.4. Lactation

IOM (1998) proposed an additional intake of 125 mg/day for lactating women aged 14–50 years, considering an average breast milk production of 0.78 L/day (Hofvander et al., 1982; Butte et al.,

1984; Chandra, 1984; Neville et al., 1988; Allen et al., 1991) and an average choline concentration of breast milk of about 160 mg/L.

An overview of DRVs for choline for infants, children, adults, pregnant or lactating women is presented in Table 1.

**Table 1:** Dietary Reference Values for choline for infants, children, adults, pregnant or lactating women

	IOM (1998) <sup>(a)</sup>
<b>Age</b> (months)	7–12
<b>All</b> (mg/day)	150
<b>Age</b> (years)	1–3
<b>All</b> (mg/day)	200
<b>Age</b> (years)	4–8
<b>All</b> (mg/day)	250
<b>Age</b> (years)	9–13
<b>All</b> (mg/day)	375
<b>Age</b> (years)	14–18
<b>Boys</b> (mg/day)	550
<b>Girls</b> (mg/day)	400
<b>Age</b> (years)	≥ 19
<b>Men</b> (mg/day)	550
<b>Women</b> (mg/day)	425
<b>Pregnancy</b> (mg/day)	450
<b>Lactation</b> (mg/day)	550

(a): Adequate Intakes. IOM: U.S. Institute of Medicine of the National Academy of Sciences.

## 5. Criteria (endpoints) on which to base Dietary Reference Values

### 5.1. Indicators of choline requirement

Plasma choline concentration may increase when intake is increased, and decreases by up to 50% when dietary intake is severely restricted (Zeisel et al., 1991; Savendahl et al., 1997) (Section 2.4.1). However, the plasma choline concentration of healthy subjects is determined not only by diet, but also by endogenous choline synthesis, potential release of choline from tissue phospholipids, microbial metabolism of dietary choline in the gut and degradation of choline via betaine (Section 2). The result of these different influences on plasma choline concentration is unpredictable. As indicated in Section 2.4.5, the Panel concludes that the available data do not allow concluding on a dose–response relationship between choline intake or choline status and plasma choline concentration. The Panel also concludes that plasma concentrations of choline, PC, betaine, DMG, tHcy or TMAO, erythrocyte PC concentration, or urinary betaine and TTMA excretion cannot be used to set DRVs for dietary choline (Section 2.4.5).

#### 5.1.1. Adults

Zeisel and co-workers performed 11 choline depletion/repletion studies in different groups of both women and men that all followed a similar design. For this reason, the characteristics of these studies are summarised below, while detailed information is available in Appendix D.

##### 5.1.1.1. Study goals

The goals differed between the studies. The first study evaluated the changes in choline status and liver function of healthy humans fed a choline-deficient diet (Zeisel et al., 1991). Another study assessed whether choline deficiency decreases the capacity to methylate homocysteine (da Costa et al., 2005) (Section 2.2.2.1). One study investigated the influence of genetic variants of folate metabolism on susceptibility to choline deficiency symptoms (Kohlmeier et al., 2005). Another assessed whether SNPs in genes coding for enzymes involved in choline metabolism influence the dietary requirement for choline and whether choline deficiency is associated with apoptosis and DNA damage



(da Costa et al., 2006b). One study investigated the influence of sex and menopausal status on dietary requirement of choline (Fischer et al., 2007). Another investigated the influence of genetic polymorphisms in *PEMT*, *MTHFD1*, *CHDH* on susceptibility for organ dysfunction in choline deficiency (Niculescu et al., 2007). One study estimated whether the risk for choline deficiency induced organ dysfunction in premenopausal women is dependent on the number of variant *PEMT* rs12325817 alleles in premenopausal women and whether oestrogen can decrease the risk in postmenopausal women (Fischer et al., 2010a). One study assessed whether metabolomic profiling of plasma can predict organ dysfunction in choline deficiency (Sha et al., 2010). One study investigated how diet and choline deficiency influence the human gastrointestinal tract microbiome and the development of liver steatosis (Spencer et al., 2011). One study assessed whether plasma PC-DHA concentration is a non-invasive marker for liver *PEMT* activity (da Costa et al., 2011). Finally, one study identified effect alleles in a number of SNPs of genes known to be of influence on the dietary requirement of choline (da Costa et al., 2014). Characteristics and outcomes of these 11 studies are compiled in Appendix D.

### 5.1.1.2. Study design

The design was similar in all studies (Appendix D), and was the following: a 7–10 day baseline diet, followed by a 42-day choline depletion diet, and then a choline-repletion diet (3–40 days). During the 10-day baseline diet, the subjects received normal foods providing 550 mg choline and 50 mg betaine/70-kg bw per day. During the choline-depletion diet, the subjects received foods providing < 50 mg choline and 6 mg betaine/70-kg bw per day for up to 42 days (with or without a folic acid supplement (100 or 400 µg/day according to study objective)), or until they were deemed choline-deficient and/or developed signs of organ dysfunction. In some studies, the participants were randomised into a depletion group and a control group that continued on the baseline diet. More details on the design per study are provided in Appendix D.

Muscle and liver dysfunction associated with choline deficiency was defined by the authors as a fivefold or greater increase in serum CK activity, a 1.5-fold or greater increase in AST, ALT,  $\gamma$ -glutamyltransferase (GGT) or lactate dehydrogenase (LDH), and/or a 28% or greater increase in liver fat content measured by computerised tomography (CT) or magnetic resonance imaging (MRI) compared with baseline and, depending on the study, estimated on day 21 and 42 of depletion. The same parameters were measured to assess reversion of the damage.

Those who completed the 42-day depletion phase without the development of hepatic steatosis were put on a diet providing 550 mg choline/70-kg bw per day for 3 days and then discharged. Choline-deficient subjects were put on a diet with stepwise increases in choline intake, in sequential 10-day periods of 137.5, 275, 412.5 or 550 mg choline/70-kg bw per day. Those who showed signs of organ damage with increases of CK activity > 10,000 U/L were immediately switched to the choline-repletion diet or directly to 850 mg choline/70-kg bw per day or to an *ad libitum* diet. Status was monitored regularly using blood and urine samples (at screening, day 1, at the end of each dietary phase, and every 3–4 days during the intervention).

### 5.1.1.3. Number of subjects and choline intake

The number of subjects ranged from 8 to 72 per study, and some of the subjects participated in several studies (Appendix D). The total number of subjects (in all studies) investigated is not quite clear, because subjects recruited in 2001 and 2007 (approximately 150–160) were investigated in different studies. The susceptibility for organ dysfunction by choline depletion was significantly influenced by polymorphisms of the *MTHFD1* (e.g. rs2236225), *CHDH*, *CHK*, *CLCA441* and *PEMT* (e.g. rs12325817) genes (Section 2.5), by being male or postmenopausal and not receiving oestrogen therapy. Folic acid supplementation (400 µg/day) did not prevent the development of organ dysfunction during choline depletion (Kohlmeier et al., 2005).

In the depletion/repletion study that investigated the influence of sex and menopausal status on choline requirement (Fischer et al., 2007) in 57 healthy adult subjects (26 males, 16 premenopausal and 15 postmenopausal women), aged 18–70 years, 20 out of 26 (77%) men developed choline deficiency signs, six already in the baseline phase with 550 mg choline/70-kg bw per day. In this study, 12 out of 15 (80%) postmenopausal women and 7 out of 16 (44%) premenopausal women developed choline deficiency signs on the low-choline diet. In total, 39 out of 57 (68%) male or female subjects developed signs of choline deficiency. In the same study, the authors also looked for differences in clinical chemistry data between subjects who developed choline deficiency and subjects who did not (apart from the parameters used to define choline deficiency-related organ dysfunction). Between sexes and life-stage groups, there were no significant differences in plasma concentrations of free

choline, betaine, DMG, tHcy, which all decreased upon depletion, and of SAM and SAH, which did not change. Plasma PC concentrations, however, decreased only in subjects who developed organ dysfunction.

The amount of choline needed to replete subjects with signs of organ dysfunction differed between subjects (Fischer et al., 2007) as shown in Table 2. In all the other studies mentioned in Appendix D, this was not reported. Disregarding missing data as well as sex differences because the numbers are too small, 10 out of 39 choline-deficient subjects were repleted with 137.5 mg/70-kg bw per day, three with 275, five with 412.5, and 13 needed 550 or more than 550 mg/70-kg bw per day (or an *ad libitum* diet) including the six men with signs of choline deficiency already on the baseline diet with 550 mg choline/70-kg of bw, while the data from eight subjects were completely missing.

**Table 2:** Amount of choline needed to replete subjects after experimental choline depletion (Fischer et al., 2007)

Study subjects	n	No signs of choline deficiency with low-choline diet	Signs of choline deficiency, with choline intake (mg/70 bw per day) of		Choline needed for repletion, total mg/70-kg bw per day				Missing data for repletion
			n	550 mg, n	50 mg, n	137.5, n	275, n	412.5, n	
Men	26	6	6*	14	6	2	3	7*	2
Premenopausal women	16	9	–	7	1	–	–	2	4
Postmenopausal women	15	3	–	12	3	1	2	4	2
Total	57	18	6*	33	10	3	5	13	8

n: number of subjects; bw: body weight.

\*Six men showed already signs of choline deficiency with 550 mg choline/70-kg bw per day and consequently needed more than that amount for repletion.

Out of 25 subjects<sup>12</sup> who showed signs of choline deficiency after experimental choline depletion and for whom the amount of choline needed to replete them was available, the Panel notes that 18, i.e. about 70%, needed up to about 400 mg choline/70-kg bw per day for repletion. The Panel also notes that this percentage decreased to 58% when the six men with signs of choline deficiency already during the baseline period with 550 mg choline/day (and therefore presumably with a higher choline requirement) were taken into account (Fischer et al., 2007). The Panel did not consider the 18 individuals who did not show signs of choline deficiency with 50 mg/70-kg bw per day. It is not known if they would have developed signs of choline deficiency with a longer period of choline depletion (> 6 weeks). The Panel notes that data are missing for the precise amount of choline needed for repletion in eight subjects.

The Panel notes that the subjects of this trial (Fischer et al., 2007) were classified according to polymorphisms in genes coding for *PEMT*, *CHDH*, *BHMT* (da Costa et al., 2006a) and for *MTFHR*, *MTFHD1* and the reduced folate carrier 1 (*RFC1*) (Kohlmeier et al., 2005) (Appendices C and D). The susceptibility to develop organ dysfunction on the low-choline diet was significantly increased ( $p = 0.002$ , odds ratio (OR): 25; 95% CI: 2–256) (18 out of 23 carriers of the C allele) in women carriers of the *PEMT* promoter SNP rs12325817 (–744 G→C), and specifically in postmenopausal women ( $p = 0.03$ , OR: 42; 95% CI: 1–1,348). In contrast, being a carrier of the *CHDH* gene SNP rs9001, +318 A→C had a protective effect on the susceptibility to develop organ dysfunction ( $p = 0.03$ , OR: 0.2; 95% CI: 0.05–0.7), while the *CHDH* SNP rs12676 (+432 G→T) did not, except in premenopausal women. The SNPs *PEMT* rs7946 (+5465 G→A) and *BHMT* rs3733890 (+742 G→A) were not associated with susceptibility to organ dysfunction on a low-choline diet. Only the *MTHFD1* SNP (1958G→A) rs2236225 carriership increased the susceptibility to develop signs of choline deficiency when the choline intake was low, and that only in premenopausal women (OR: 85, 95% CI: 3–2,418), and this susceptibility was attenuated by folate supplementation.

There are indications that choline deficiency during depletion/repletion studies (da Costa et al., 2006a; Niculescu et al., 2007) (Appendix D) may increase cell apoptosis and induce DNA damage

<sup>12</sup> i.e. 10 + 3 + 5 + 13–6, indicated in Table 2.

(assessed *ex vivo/in vitro*), for which the carriers of certain polymorphisms of *PEMT* and *MTHFD1* were more susceptible (Section 2.5). The Panel considers that the significance of these studies is unclear.

There are also indications (Appendix D) that metabolomic profiling of the plasma of subjects on baseline diet can predict susceptibility to develop organ dysfunction when deprived of dietary choline (Sha et al., 2010) and that host factors and the gut microbiota (Spencer et al., 2011) both respond to dietary choline intake and choline deficiency (Section 2.2.2.1).

#### 5.1.1.4. Summary

Eleven available depletion/repletion studies in adults have demonstrated that dietary choline can become insufficient, e.g. within 6 weeks of a depletion phase with  $\leq 50$  mg choline/70-kg bw per day (Appendix D). Only one of these studies reported the amount of choline needed to replete subjects with signs of organ dysfunction (Fischer et al., 2007).

The Panel notes that experimental dietary depletion of choline led, in most (70–80%) of the male and postmenopausal female subjects, to signs of organ dysfunction involving liver and muscle, but only in 44% of premenopausal women (Fischer et al., 2007). These signs can be mild with biochemical alterations only or can be severe with liver steatosis and muscle function impairment developing rapidly. The susceptibility to develop organ dysfunction differs between subjects and is influenced by genetics, sex, possibly the intestinal microbiome, and hormonal status (Section 2).

In addition, it is not known if the 18 subjects who have not developed signs of organ dysfunction within 6 weeks would have done so in the long term, when their endogenous choline (PC) synthesis would become insufficient (Fischer et al., 2007). It is not known, but can be assumed, that the factors that have an impact on the development of organ dysfunction also determine the amount of choline needed to replete the body and reverse the signs of organ dysfunction and the requirement for dietary choline.

According to the study by Fischer et al. (2007) described above, this requirement for dietary choline in adults lies between about 130 and 500 mg choline/day, with most subjects needing more than 130 mg/day and some needing 500 mg/day or more (Table 2). From the 39 subjects who became deficient either with 550 or with 50 mg choline/70-kg bw per day, the data from 14<sup>13</sup> are missing. From the remaining 25, ten needed 137.5, three 275, five 412.5 and seven 550 mg choline/70-kg bw per day or more. An intake of 412.5 mg choline/70-kg bw per day (i.e. 5.9 mg/kg bw per day) was sufficient to replete 18 of 25 deficient subjects, that is about 70% or two-thirds.

The Panel considers that reliable markers of intake and status are not available (Section 2.4) and that the study by Fischer et al. (2007) is too small and insufficient to draw firm conclusions on the Average Requirement (AR) for dietary choline in adults. However, as supportive evidence, it may contribute to inform an AI that covers most of the population.

#### 5.1.2. Infants and children

The Panel is unaware of any data in infants aged 7–11 months and children on indicators of choline requirement.

#### 5.1.3. Pregnancy and lactation

The Panel considered whether the calculation of choline transfer from the mother to the fetus and of choline accretion in the fetus and placenta during pregnancy could be used to calculate the additional need for dietary choline during pregnancy. However, a review of the available evidence (Sections 2.3.3 and 2.3.4) showed that this was not feasible due to a lack of data.

The Panel then considered the available intervention studies on choline supplementation in pregnant women in the second half of pregnancy, and in lactating women. Although none of the biomarkers in plasma, urine or erythrocyte previously reviewed by the Panel are suitable biomarkers to set DRVs for choline (Section 2.4.5), the Panel considers that they may be useful to assess potential changes in choline metabolism in intervention studies in pregnant or lactating women.

##### 5.1.3.1. Effect of total intake of choline in pregnant (vs non-pregnant) women and the offspring

As described already in Sections 2.3.6.1 and 2.4.1.2, Yan et al. (2012) reported on plasma and urine choline concentrations in 26 healthy pregnant women (third trimester) and 21 non-pregnant

<sup>13</sup> 8 + 6 (Table 2).



controls who were randomly assigned to consume either 480 or 930 mg of choline/day from food and supplements for 12 weeks (or until delivery). The diet provided an average of 380 mg/day of choline (sum of all compounds), and supplemental choline was 100 or 550 mg/day (from choline chloride).<sup>14</sup> Pregnant women had higher free choline concentration in plasma and urinary excretion of choline and betaine than non-pregnant women throughout the study (Sections 2.3.6.1 and 2.4.1.2). Also, pregnant women consuming 930 mg of choline/day had higher plasma concentrations of free choline than pregnant women consuming 480 mg of choline/day. The lower circulating concentrations of choline-derived methyl-group donors (betaine, DMG and sarcosine) observed in pregnant women compared with non-pregnant women were suggestive of a greater use of these molecules in both maternal and fetal compartments (Section 2.4.1.2).

This study also provided additional results. Plasma concentrations of the three methyl-group donors (betaine, DMG and sarcosine) over the duration of the study were higher in pregnant women consuming 930 mg choline/day compared with pregnant women consuming 480 mg of choline/day ( $p < 0.016$ ,  $p < 0.012$ , and  $p < 0.07$ , respectively), but without achieving the concentrations measured in non-pregnant women consuming 480 mg choline per day. Urinary excretion of choline, betaine or DMG in pregnant women was not different between the choline intake groups. However, urinary excretion of sarcosine, methionine and Hcy were higher (46% higher,  $p = 0.029$ ; 37% higher,  $p = 0.02$ ; 45% higher,  $p = 0.06$ , respectively) in the pregnant women consuming 930 mg/day, compared with 480 mg/day. The results described above in plasma and in urine suggest that the higher choline intake (930 mg/day) was predominantly used by the pregnant women, and not excreted. However, in pregnant women, mean concentration of free choline in the placenta ( $915 \pm 231$  vs  $941 \pm 309$  nmol/g tissue) or in cord plasma ( $37.3 \pm 13$  vs  $32.5 \pm 7.5$   $\mu\text{mol/L}$ ), and anthropometric parameters or Apgar scores of the newborns did not differ between the lower and the higher choline intake groups.

#### 5.1.3.2. Effect of total intake of choline in pregnant (vs non-pregnant) women on the dynamics of choline-related metabolic pathways

As indicated previously, the PC formed in the PEMT pathway contains substantial amounts of LC-PUFAs, like DHA and ARA, while the PC formed in the CDP-choline pathway does not (Section 2.3.5).

Yan et al. (2013) investigated the effect of pregnancy on the dynamics of choline-related metabolic pathways (Figure 2, Section 2.3.5) in the same study cohort of pregnant (third trimester) and non-pregnant women investigated by Yan et al. (2012) who had received, after 6 weeks, 100 mg (of the 480 mg/day choline) and 200 mg (of the 930 mg/day choline) as deuterated choline (methyl-D<sub>5</sub>-choline). In pregnant women (compared with non-pregnant women), the total plasma PC pool was about 50% greater (Yan et al., 2013).

With regard to the CDP-pathway, the analysis of the different isotopomers of deuterated choline, betaine and PC in plasma showed that in pregnant women (compared with non-pregnant women), dietary choline was used more for PC production via the CDP-choline pathway than oxidised to betaine. The higher choline intake (930 mg choline/day) in pregnant women restored the distribution of dietary choline between PC synthesis via the CDP-choline pathway vs oxidation to betaine, to the levels observed in non-pregnant women consuming 480 mg choline/day. With regard to PEMT pathway, the analysis of the different isotopomers also showed that, in pregnant women (compared with non-pregnant women), PC produced via PEMT is more catabolised to free choline (and this may contribute to explain the rise in plasma choline in pregnancy), which is preferentially transferred to the fetus. The higher choline intake (930 mg choline/day) enhanced the PEMT-mediated PC synthesis relative to the CDP-choline pathway, compared to pregnant women consuming 480 mg choline/day.

West et al. (2013) investigated the effect of different choline intakes on choline-related lipid metabolism in a separate analysis of the same study cohort of pregnant (third trimester) and non-pregnant women investigated by Yan et al. (2012). At baseline, pregnant women had a greater proportion of PC-DHA (% of total fatty acids) in both plasma ( $p = 0.01$ ) and erythrocytes ( $p = 0.001$ ) than non-pregnant women. The higher choline intake (930 mg/day) did not affect the proportion of PC-DHA in erythrocytes in pregnant women compared with an intake of 480 mg/day (whereas this was the case in non-pregnant women, as described in Section 2.3.5.1). However, the higher choline intake (930 mg/day) lowered the proportion of PC-ARA in erythrocytes in pregnant women ( $p = 0.02$ ),

<sup>14</sup> In addition to the strictly controlled diet, all subjects received 600  $\mu\text{g}$  folic acid, 2.6  $\mu\text{g}$  cobalamin, 1.9 mg vitamin B6 and 200 mg DHA per day.

compared with an intake of 480 mg/day. The PC:PE ratio (Section 2.3.5.1) in plasma and erythrocytes was not influenced by choline intake in pregnant or non-pregnant women.

#### 5.1.3.3. *Ex vivo* studies in placental samples

From 24 subjects from the study by Yan et al. (2012) (12 each from the two choline groups), placental tissue, cord blood leukocytes and maternal fasting venous blood at delivery were investigated *ex vivo* by Jiang et al. (2012, 2013). In the group that consumed 930 mg/day choline compared with the group that consumed 480 mg/day choline, the authors found that: (i) placental global DNA methylation, histone methylation and the expression of a histone methyltransferase were higher; (ii) placental methylation of the promoters of two cortisol-regulating genes, corticotropin releasing hormone (*CRH*) and glucocorticoid receptor (*NR3C1*) was higher; (iii) placental *CRH* transcript abundance was lower (about 40%, read on figure, concentration of the protein was not reported); (iv) methylation of the *CRH* and *NR3C1* promoter in cord blood leukocytes was lower; (v) the maternal blood concentration of the protein antiangiogenic factor fms-like tyrosine kinase (*sFLT1*) at delivery was lower (by about 30%, estimated from the figure); (vi) placental *sFLT1* mRNA abundance was lower (by about 30%, estimated from the figure, concentration of the protein was not reported).

#### 5.1.3.4. Effect of total intake of choline on maternal plasma and breast milk during lactation

The RCT by Fischer et al. (2010b) (Sections 2.3.3, 2.3.6.3, 2.4.1.2, and 5.1.1.1) demonstrated that total intake of choline (from foods and supplements) is positively associated with the concentration of free choline and choline-compounds in plasma of these lactating women (Section 2.4.1.2) and in breast milk (Section 2.3.6.3). This study also showed that supplemental choline (750 mg/day choline, in addition to a mean dietary choline intake of about 350 mg/day) compared with placebo increased the mean concentration of free choline in plasma (Section 2.4.1.2) and in breast milk (Section 2.3.6.3).

In the previously described controlled feeding study by Davenport et al. (2015) (Sections 2.3.3, 2.3.6 and 2.4.1.2), lactating and control non-lactating women (the latter from the study by Yan et al. (2012)) were randomised to consume 480 mg choline/day or 930 mg choline/day from food and supplements<sup>7</sup> for 10–12 weeks, and they all received, during the last 4–6 weeks, 20% of the total intake of choline as deuterium labelled choline. Lactating (vs control) women showed a statistically lower expression of three of the five genes investigated that code for enzymes/receptor involved in choline metabolism, in leukocytes at baseline (mRNA abundance,  $p \leq 0.05$ ). They also showed a higher plasma free choline concentration (Section 2.4.1.2) and lower urinary excretion of choline metabolites (Section 2.3.6.1.2) throughout the study period. Lactating (vs control) women tended to have a decreased oxidation of choline to betaine (Figure 2, Section 2.3.5), which would allow an increase in the supply of intact choline to the mammary epithelium. The higher choline intake during lactation (930 mg/day compared to 480 mg/day) significantly increased the concentration of total choline in breast milk and increased the supply of PEMT-derived choline metabolites in breast milk (Section 2.3.6.3), as well as in blood.

#### 5.1.3.5. Conclusion on pregnancy and lactation

In pregnant women (compared to non-pregnant women) (Yan et al., 2012, 2013; West et al., 2013), the available studies:

- show increased urinary losses of choline and betaine;
- suggest a greater use of choline-derived methyl-group donors (DMG, betaine and sarcosine) in both maternal and fetal compartments;
- suggest an enhanced PEMT activity to facilitate the transfer of LC-PUFA to the fetus via PC in lipoproteins.

These studies on choline supplementation also suggest that a choline intake of 930 mg/day (from food and supplements) in pregnant women (from the 27th week of gestation):

- increases (compared to 480 mg/day) maternal plasma choline concentration;
- increases maternal plasma concentrations of the three methyl-group donors (DMG, betaine and sarcosine) compared with pregnant women consuming 480 mg/day, but without achieving the concentrations measured in non-pregnant women consuming 480 mg/day;

- restored the distribution of dietary choline between PC synthesis via the CDP-choline pathway vs oxidation to betaine to the levels observed in non-pregnant women consuming 480 mg choline/day;
- enhanced (compared to 480 mg/day) the PEMT-mediated PC synthesis vs the CDP-choline pathway-mediated PC synthesis;
- had no impact (compared to 480 mg/day) on maternal urinary excretion of choline and betaine, placental choline concentration, and cord plasma choline concentration.

These results may indicate a higher choline requirement in pregnancy than in non-pregnant women, which would have to be supplied by additional dietary choline.

In lactating women, the available studies on choline supplementation on women either supplemented from the 18th gestational week to 45 days post partum (Fischer et al., 2010b) or recruited at 5 weeks post partum (Davenport et al., 2015), suggest that increased maternal choline intake enhances the concentration of total choline in breast milk and increased the supply of PEMT-derived choline metabolites in breast milk. As PEMT generates PC molecules enriched in DHA, the supply of DHA from the lactating women to the infant might be facilitated. However, the fatty acid composition of breast milk was not measured in these studies.

The Panel notes that no maternal clinical signs of choline deficiency (as described in Sections 2.2.2.1 and 5.1.1.4) or no adverse outcomes in the offspring were reported in these studies with a total intake of choline from foods and supplements of 480 mg choline/day in pregnant women, or of about 350–480 mg choline/day in lactating women.

The Panel notes that these studies used high choline intakes (930 vs 480 mg/day from foods and supplements in pregnant and lactating women in one study; about 1,100 mg/day from food and supplements vs about 350 mg/day from foods in lactating women in another study). The Panel also notes that the interpretation of the biochemical outcomes investigated is difficult with the aim of defining choline insufficiency/adequacy in pregnancy.

The Panel notes that the *ex vivo* studies suggest that different maternal choline intakes during pregnancy may induce epigenetic modifications of genes, and changes in genes involved in hormonal and vascular physiology. However, such changes are difficult to interpret and further research is required.

The Panel concludes that calculation of the additional need for dietary choline during pregnancy based on a calculation of choline transfer from the mother to the fetus and choline accretion in the fetus and placenta during the duration of pregnancy is not feasible due to a lack of data (Sections 2.3.3 and 2.3.4). The Panel concludes that, taken together, the studies on choline supplementation provide evidence that pregnant or lactating women may need more choline than non-pregnant non-lactating women. However, the data are not sufficient to allow an estimate of the additional requirement for dietary choline in pregnant or lactating women (above that of non-pregnant non-lactating women). The Panel considers, however, that the additional intake of choline required to compensate for the amount of total choline secreted in breast milk during the first 6 months of exclusive breastfeeding (Section 2.3.6.3) can be calculated.

## 5.2. Choline intake and health consequences

Since the report by SCF (1993), more data have become available on the relationship between choline intake and NAFLD, CVD, different types of cancer, NTD and cognition. A comprehensive search of the published literature, without time limit, was performed in August 2012 as preparatory work to this Opinion in order to identify relevant health outcomes possibly associated with choline intake through diet or supplementation, and which may inform the setting of DRVs for choline (El-Sohemy et al., 2012). The main results of the preparatory work, together with new evidence from studies subsequently published (in Pubmed) until November 2015 are summarised below.

Of the available RCTs investigating the health effects of choline, the results only of one RCT was considered in this section, which reported dietary choline intake in addition to choline supplements. The relationship between choline intake and chronic disease outcomes has been investigated mainly in observational (prospective cohort, case-control) studies, where a positive, an inverse, or a lack of an association between choline intake and disease outcomes might be confounded by uncertainties inherent to the methodology used for the assessment of choline intakes, and by the effect of other dietary, lifestyle or undefined factors on the disease outcomes investigated. Taking into account the uncertainty about the relationship between choline intake and biomarkers (Section 2.4), the Panel only considered observational studies that include an assessment of choline intake, whereas studies on the

relationship of plasma choline concentrations (or those of choline compounds) and health outcomes with no quantitative data on choline intake (Wang et al., 2011) are not described below. In observational studies, habitual dietary choline intake was generally estimated using a FFQ (filled-in either once at baseline or at several time points, in prospective cohort studies) and composition data from the USDA database (Section 3) and/or from the literature (Zeisel et al., 2003). For some observational studies, choline intake from supplements was also assessed.

### 5.2.1. Non-alcoholic fatty liver disease

Dietary deficiency of choline can cause fatty liver (hepatic steatosis), which can result in NAFLD (Section 2.2.2.1), which can be of different aetiologies and is the most common chronic liver disease in developed countries. It is often associated with insulin resistance and dyslipidaemia, is a risk factor for CVD and may progress to irreversible liver damage and liver cancer (Corbin et al., 2013; Lazo et al., 2013; Byrne and Targher, 2014).

In two population-based prospective cohorts, Yu et al. (2014) investigated the association between habitual dietary choline intake and risk of NAFLD in 56,195 women (recruited in 1997–2000 and followed-up through 2004–2007) and men (recruited in 2002–2006 and followed-up through 2008–2011), aged 40–75 years and free of hepatitis at baseline. NAFLD was diagnosed by sonography (self-report). Mean daily choline intake was 412 mg (women) and 452 mg (men) in the highest quintile, and 179 mg (women) and 199 mg (men) in the lowest quintile. After adjustment for potential confounders,<sup>15</sup> women and men in the highest quintile had a significantly lower risk of NAFLD than those in the lowest quintile, but not after further adjustments. In stratified analysis, the highest quintile of choline intake remained inversely associated with risk of NAFLD compared with the lowest quintile (OR: 0.72; 95% CI: 0.57–0.91, *p* trend = 0.007) only in women with a body mass index (BMI) < 25 kg/m<sup>2</sup> (but not in women with a BMI ≥ 25 kg/m<sup>2</sup>).

The Panel notes that, in one prospective cohort study, a lower choline intake was associated with a higher risk of developing NAFLD in normal-weight women in adjusted stratified analysis. The Panel concludes that the data on choline intake and risk of NAFLD are limited and cannot be used to derive DRVs for choline.

### 5.2.2. Cardiovascular disease

A prospective cohort study, with an average follow-up of 8.1 years, investigated the association between habitual dietary intake of choline and risk of CVD in 16,165 postmenopausal women aged 49–70 years and without prior CVD at baseline (Dalmeijer et al., 2008). After adjustment for potential confounders, comparing the highest quartile of choline intake (> 329 mg/day) with the lowest (< 266 mg/day) did not show a significant relationship between choline intake and risk of total CVD, coronary heart disease (CHD) or cerebrovascular accidents.

A prospective cohort study, with an average follow-up of 14 years, investigated the association between habitual dietary intake of choline and risk of CHD, in 14,430 men and women without prior CHD at baseline (mean age at baseline: about 54 years) (Bidulescu et al., 2007). After adjustment for potential confounders, comparing the highest quartile of choline intake (> 363 mg/day) with the lowest (< 217 mg/day) did not show a significant relationship between choline intake and risk of CHD.

The Panel notes that two large prospective cohort studies on populations free of CVD at baseline did not show a significant association between choline intake and risk of CVD. The Panel concludes that the data on choline intake and risk of CVD cannot be used to derive DRVs for choline.

### 5.2.3. Cancer

Choline is a methyl-group donor involved in the folate-dependent one-carbon metabolism (Sections 2.2.1 and 2.3.5). Disturbances in this function that affect methylation or synthesis of DNA may contribute to carcinogenesis (Section 2.2.2.1).

#### 5.2.3.1. Colon/rectum

In a US prospective cohort study, Cho et al. (2007a) examined the relationship between total intake of choline (via food and supplements) and risk of colorectal adenoma in 39,246 women free of cancer

<sup>15</sup> Including age, total energy intake, education, income, physical activity, smoking, alcohol consumption, intake of protein, saturated fat, polyunsaturated fat. Further adjustments for menopause, hypertension, diabetes mellitus, gallstones, dyslipidaemia, BMI.



or polyps at baseline and who underwent at least one endoscopy in the 18 years of follow-up. After adjustment for potential confounders, a choline intake in the highest quintile (median: 383 mg/day) was associated with a higher risk of colorectal adenomas compared with the lowest quintile (median: 261 mg/day) (relative risk (RR): 1.45; 95% CI: 1.27–1.67;  $p$  trend < 0.001).

In a US prospective cohort study, Lee et al. (2010a) investigated the relationship between total intake of choline (via food and supplements) and risk of colorectal cancers (CRCs) in 47,302 men (40–75 years at baseline) free of cancer at baseline and with 18 years of follow-up. After adjustment for potential confounders, a choline intake in the highest quintile, from either food or supplements, was not associated with a higher risk of CRC compared with the lowest quintile.

In a case–control study, Lu et al. (2015) investigated the relationship between habitual dietary intake of choline and risk of CRC, in 890 cases (aged 30–75 years) diagnosed up to 3 months previously, compared with 890 age- and sex-matched controls. Choline intake (median, 25th and 75th percentiles) was higher in controls (158, 120 and 202 mg/day) than in cases (133, 100 and 176 mg/day) ( $p$  < 0.01). After adjustment for potential confounders, a choline intake in the highest quartile was inversely associated with risk of CRC compared with the lowest quartile (OR: 0.54; 95% CI: 0.37–0.80;  $p$  trend < 0.01). The Panel notes that the diet in this population provided about half of the dietary choline and folate intake, and less red meat, poultry, eggs and milk than in the USA (Cho et al., 2007a).

The Panel notes the inconsistent results from observational studies on the association between choline intake and risk of colorectal cancer.

### 5.2.3.2. Breast cancer

In a prospective cohort study with a follow-up of 12 years, Cho et al. (2007b) examined the relationship between total intake of choline (via food and supplements) and risk of breast cancer in 90,663 premenopausal women, aged 26–46 years, and free of cancer at baseline. Median intake per quintile ranged between 263 and 397 mg/day. After adjustment for potential confounders, choline intake was not associated with breast cancer risk.

In a prospective cohort study, Cho et al. (2010) investigated the relationship between habitual dietary intake of choline and risk of breast cancer in 74,584 women, who were either postmenopausal in 1984 or became postmenopausal during 20 years of follow-up (mean age of about 62 years at 10-year follow-up). Median intake per quintile ranged between 260 and 396 mg/day. After adjustment for potential confounders, choline intake was not associated with breast cancer risk.

In a population-based case–control study, Xu et al. (2009) investigated the relationship between total intake of choline (via foods and supplements) and risk of (and mortality from) breast cancer and all-cause mortality in 1,508 cases of breast cancer (diagnosed in 1996–1997 and followed through 2005) and 1,556 controls. After adjustment for age, choline intake (sum of all forms, ranging from < 123 mg/day to > 247 mg/day) was not associated with risk of breast cancer. In addition, choline intake (sum of all forms, ranging from < 142 to > 205 mg/day) was not associated with all-cause or breast cancer mortality (while an inverse significant relationship for both types of mortality was observed comparing intake of free choline above about > 57 mg/day with that < 40 mg/day).

The Panel notes that three observational studies did not show a significant association between choline intake and risk of breast cancer. The Panel concludes that the data on choline intake and risk of breast cancer cannot be used to derive DRVs for choline.

### 5.2.3.3. Other cancers (oesophageal, prostate and ovarian cancers)

In two population-based case–control studies, Ibiebele et al. (2011) evaluated the association between habitual dietary intake of choline and risk of Barrett's oesophagus (BE) and oesophageal cancers. The first study compared eligible cases ( $n$  = 367), diagnosed with BE or BE with dysplasia, with 577 controls. The second study compared eligible cases ( $n$  = 881), diagnosed with oesophageal carcinoma of different types and location, with 1,507 controls. Median intake of choline in each quartile in controls ranged between 380 and 1,171 mg/day. After adjustment for potential confounders, choline intake was not associated with risk of BE or oesophageal cancers.

In a prospective cohort study with a follow-up of 22 years, Richman et al. (2012) examined the association between total intake of choline (via foods and supplements) and risk of fatal prostate cancer in 47,896 men, aged 40–75 years, and free of cancer diagnosis at baseline. After adjustment for potential confounders, the highest quintile of choline intake (median 509 mg/day) was positively associated with risk of fatal prostate cancer (hazard ratio (HR): 1.70; 95% CI: 1.18–2.45,  $p$  trend = 0.005).



In two large prospective cohorts with a follow-up of up to 22 years, Kotsopoulos et al. (2010) investigated the relationship between total intake of choline (via foods and supplements) and risk of ovarian cancer, among 159,957 women, aged 25–55 years at enrolment. In both cohorts, choline cutpoints ranged between about 250–270 mg/day (lowest quintile) and 339–367 mg/day (highest quintile). After adjustment for potential confounders, choline intake was not associated with risk of ovarian cancer.

The Panel notes that choline intake was not associated with risk of oesophageal cancer in one reference on two case–control studies or with risk of ovarian cancer in two cohorts followed prospectively, while it was positively associated with risk of prostate cancer in one large prospective cohort study.

#### 5.2.3.4. Conclusions

The Panel concludes that the available data on associations between choline intake and cancers of various sites are either inconsistent or limited and cannot be used to derive DRVs for choline.

#### 5.2.4. Neural tube defects

In a US population-based case–control study, Shaw et al. (2004) investigated the relationship between periconceptual intake of choline and risk of NTDs, in 653 cases (liveborn, stillborn or electively terminated) identified from hospital and medical records (in 1989–1991), compared with 644 controls randomly selected from the same geographical area. Dietary choline intake of the mothers (not taking supplements with choline) in the 3 months before conception was estimated retrospectively. The authors analysed 424 FFQs from mothers of NTD cases (161 with anencephaly, 242 with spina bifida, 21 with other NTD phenotypes) and 440 FFQs of controls. After adjustments for potential confounders, a significantly decreased risk of all NTDs was found for quartiles 2–4 of periconceptual intake of choline compared to the lowest quartile (< 290 mg/day), e.g. for the fourth quartile (> 498 mg choline/day) OR: 0.49; 95% CI: 0.27–0.90.

In another US population-based case–control study, Carmichael et al. (2010) investigated the relationship between periconceptual intake of choline and risk of NTDs in 189 cases of spina bifida and 141 cases of anencephaly (liveborn, stillborn, electively terminated) identified from hospital and medical records (in 1999–2003), compared to 625 controls randomly selected from the same geographical area. Dietary choline intake of the mothers in the 2 months before/after conception was estimated retrospectively (8–10 months after delivery). After adjustments for potential confounders, periconceptual intake of choline (supplements excluded) below the 25th percentile (< 293 mg/day) and above the 75th percentile (> 506 mg/day) was not associated with a higher or lower risk for anencephaly and spina bifida, compared to a choline intake between the 25th and 75th percentiles.

Polymorphisms in genes for enzymes (CHKA, MTHFD1 and CCT) involved in choline metabolism may influence the risk of NTDs independently of maternal choline intake (Appendix C and Section 2.5), but such information is not available for the studies cited above.

The Panel notes that the association between choline intake and risk of NTDs was inconsistent in the two case–control studies available, and that such association may be influenced by the intake of other nutrients and the genotype of the mother. The Panel concludes that the data on choline intake and risk of NTDs cannot be used to derive DRVs for choline.

#### 5.2.5. Cognition

The only RCT, then the prospective observational studies (first in adults, then in children) are described below.

In a double-blind RCT, Cheatham et al. (2012) investigated the relationship between maternal PC supplementation during and after pregnancy (in women that, for most of them, had been investigated by Fischer et al. (2010b)) and several measures of cognition in the infants. From 18 weeks of gestation to 90 days post partum, 140 healthy women (Section 2.3.3, 2.3.6.3, 2.4.1.2, 2.5.1, 5.1.3) received either 750 mg/day of choline (as PC, n = 49 included in the analysis) or a placebo (n = 50 included in the analysis), in addition to a diet providing a mean of about 360 mg/day choline (assessed at 30 weeks of gestation and 45 days post partum). Infants (n = 99) were breast-fed for at least 45 days, and were assessed for short-term visuospatial memory (with a Delayed Response Task), long-term episodic memory (with a deferred imitation task), language development (with the Mac-Arthur Bates Short Form Vocabulary Checklist) and global development (with the Mullen Scales of Early Learning) at 10 and 12 months of age. There were no significant differences between the groups on any of the cognitive assessments at either age.

In a prospective cohort study, Poly et al. (2011) investigated the association between habitual dietary intake of choline and performance at a neuropsychological test battery or brain morphology, assessed by MRI, in 1,391 men and women (aged 36–83 years) without dementia at baseline. Choline intake was estimated in 1991–1995 with the Harvard FFQ, and again in 1998–2001 when a neuropsychological test battery and a brain MRI scan were also administered. Factor analysis was used to identify four cognitive factors (verbal memory, visual memory, verbal learning and executive function) from the numerous individual neuropsychological tests. Mean choline intake was about 322 mg/day in both periods. After adjustment for potential confounders, performance on the verbal memory and visual memory factors were significantly better with higher choline intake in 1998–2001 ( $p < 0.01$ ) but there were no significant effects for verbal learning and executive function. No significant association between choline intake (either period) and total cranium brain volume was found.

In a prospective prebirth cohort in 2,128 pregnant women included at less than 22 weeks of gestation, Villamor et al. (2012) investigated the relationship between maternal total intake of choline (via foods and supplements), assessed with an FFQ during the first and second trimesters of pregnancy, and performance on cognitive tests in their children ( $n = 1,210$ ) at 3 years of age. The cognitive tests included the Peabody Picture Vocabulary Test III and the Wide Range Assessment of Visual Motor Abilities. Maternal intake of choline (mean  $\pm$  SD) was  $332 \pm 63$  and  $325 \pm 64$  mg/day in the first and second trimesters, respectively. There was no association between maternal choline intake at either trimester and cognitive outcomes, after adjustment for potential confounders.

However, in this same cohort, Boeke et al. (2013) assessed 890 children with complete data at the age of 7 years for visual memory (measured with the Wide Range Assessment of Memory and Learning Second Edition (WRAML2), Design and Picture Memory subtests) and both verbal and non-verbal intelligence, measured with the Kaufmann Brief Intelligence Test, Second Edition (KBIT-2)). The top quartile of second trimester maternal choline intake (median (range): 392 (364–806) mg/day) was significantly associated with a WRAML2 score 1.4 points higher (95% CI: 0.5–2.4,  $p$  trend = 0.003) than the bottom quartile (median (range): 260 (141–288) mg/day), after adjustment for potential confounders. The association was not statistically significant for the first trimester maternal choline intake. Comparing the top quartile of second trimester maternal intake with the first quartile, the effect estimate for the child non-verbal KBIT-2 score was 3.5 (95% CI: 0.1–6.9;  $p$  trend = 0.06).

The Panel notes that one RCT found no difference in four cognitive parameters investigated in infants, at 10 and 12 months of age, whose mothers had consumed 750 mg/day choline or placebo in addition to their choline intake from the diet during the third trimester. The Panel also notes that available data on the relationship between choline intake and cognition in adults are limited. The Panel also notes the discrepancy in the results of a prospective cohort study, investigating the relationship between maternal choline intake during the first and second trimesters of pregnancy and cognitive outcomes in the children, when these children were aged three or 7 years. The Panel considers that this might suggest that to investigate the effects of prenatal choline supply on visual memory of the children, long-term observations are needed, and that the available evidence is insufficient to demonstrate a causal relationship. The Panel concludes that the data on choline intake and cognition cannot be used to derive DRVs for choline.

#### 5.2.6. Conclusion on choline intake and health consequences

In studies pointing to an association of higher choline intake with a reduced risk for a certain outcome (i.e. risk of liver steatosis or of NTDs, one study each), the beneficial effect was associated with choline intakes between about 400 and 500 mg/day. However, one adverse health outcome (higher risk of prostate cancer in one study) was associated with similar choline intakes (Section 5.2). The Panel concludes that the data on choline intake and health outcomes are either limited or inconsistent or do not show a significant association, and, therefore, cannot be used to derive DRVs for choline. There is a lack of data on choline intake in infants in the second half year of life and children and on associations between choline intake and health outcomes in children that could be used to set requirement for choline in these age groups.

## 6. Data on which to base dietary reference values

### 6.1. Adults

Mean observed intakes of healthy adults of all ages in Europe ranged from about 270 to 470 mg choline/day (Section 3.2.1), and the midpoint of this range is around 370 mg/day.

The Panel notes that choline depletion/repletion studies (Section 5.1.1) indicate large variability in dietary choline requirement. The Panel also notes that the variability in choline requirement due to differences in sex, polymorphisms of genes coding for enzymes involved in choline and folate metabolism, nutritional and hormonal status, and likely the composition of the gut microbiome, pose a difficulty for dose-finding studies in a sufficiently large sample of the population (Section 2). The Panel concludes that choline depletion/repletion studies do not provide sufficiently precise data to calculate Average Requirements (ARs) and PRIs for dietary choline.

The Panel also notes that there is only one depletion/repletion study that reports the choline amounts that were needed/sufficient to reverse the signs of choline deficiency in a small number of subjects (Fischer et al., 2007). In this study, out of 25 subjects who showed signs of choline deficiency after experimental choline depletion and for whom the amount of choline needed to replete them was available, about two-thirds (or about 70%) of subjects needed up to about 400 mg choline/70-kg bw per day for repletion (Table 2, Section 5.1.2).

Finally, the Panel chose to set an AI for choline for adults based on data on observed mean intakes in healthy populations, investigated in 12 national surveys undertaken in nine countries in the EU between 2000 and 2011 (Section 3.2.1), and in consideration of the amount of choline needed to replete about two-thirds (or about 70%) of choline-depleted subjects who showed signs of organ dysfunction and for whom data on the amount of choline needed for repletion were available (Fischer et al., 2007). The Panel is aware of the inherent uncertainty of the chosen value. However, assuming that the choline requirement of the 18 subjects of this study who did not show signs of choline deficiency after a restriction of the choline intake to 50 mg/70-kg bw per day for 6 weeks, will also be covered by an intake of 400 mg/day, the Panel considers this choice of 400 mg/day to be a safe and conservative approach.

Although premenopausal women may have a lower requirement for dietary choline than postmenopausal women, in connection with a potential stimulation of the PEMT pathway by oestrogen, the Panel is not aware of quantitative data with regard to the enhanced activity of the PEMT. Although ranges of estimated mean observed choline intake in healthy populations in the EU are slightly lower in women than men (Section 3.2.1), and considering that the data from the one depletion/repletion study (Fischer et al., 2007) are insufficient to conclude on sex-specific DRVs, the Panel considered it unnecessary to give sex-specific AIs for adults.

The Panel proposes an AI of 400 mg/day for all adults.

## 6.2. Infants

Considering that there is no evidence for an insufficient choline intake of fully breast-fed infants during the first 6 months of life, the amount of choline provided in human milk is considered to be adequate. Considering a choline concentration of 145 mg/L (average of two studies on full-term infants) and assuming a mean milk transfer of 0.8 L/day during the first 6 months of lactation in exclusively breastfeeding women (Butte et al., 2002; FAO/WHO/UNU, 2004; EFSA NDA Panel, 2009), the estimated choline intake of fully breast-fed infants during the first 6 months of life would be 116 mg/day, rounded up to 120 mg/day (Section 2.3.6.3).

In order to estimate the AI of infants aged 7–11 months by upwards extrapolation from the calculated choline intake for exclusively breast-fed infants from birth to 6 months, allometric scaling was applied. The Panel calculated averages of the median weights of male and female infants, aged 3 months (6.1 kg) and 9 months (8.6 kg); the median weight-for-age data came from the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006).

$$AI_{\text{infants 7–11 months}} = \text{choline intake}_{\text{infants 0–6 months}} \times \left( \frac{\text{weight}_{\text{infants 7–11 months}}}{\text{weight}_{\text{infants 0–6 months}}} \right)^{0.75}$$

This calculation yields a value of 155, which gives an AI of 160 mg/day after rounding (Table 3).

**Table 3:** Reference body weights and Adequate Intake (AI) of choline for infants aged 7–11 months

Age	Reference body weight (kg)	AI (mg/day)
7–11 months	8.6 <sup>(a)</sup>	160

(a): Average of the median weight-for-age of male or female infants, respectively, aged 9 months according to the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006).

### 6.3. Children

The Panel recognises the limited number of data on age-specific choline intake in European children and uncertainty surrounding these data (Section 3.2). The Panel chose to derive AIs for all children by downwards extrapolation from the AI for adults (400 mg/day) (Section 6.1), taking into account that this AI for adults was based on data on observed intakes in the EU, and the amounts of choline needed to replete about two-thirds (or about 70%) of choline-depleted adults who had developed signs of organ dysfunction and for whom data on choline amounts needed for repletion were available. This downwards extrapolation was carried out based on reference body weights using allometric scaling with age-dependent growth factors, and applying the 0.75 power of body mass to correct for differences in the metabolically active body mass of subjects of different sizes. Although it is not known if the choline requirement is related to energy metabolism, the Panel considers that allometric scaling, which results in a higher percentage of the adult AI than when the actual body weight is used, is justified to cover the need for choline in the development of organs and their composition.

No data are available that would justify different AIs for boys and girls.

The AIs were calculated by using the following equation

$$AI_{\text{child}} = AI_{\text{adults}} \times \left( \frac{\text{weight}_{\text{child}}}{\text{weight}_{\text{adults}}} \right)^{0.75} \times (1 + \text{growth factor})$$

For the calculations (Table 4), median body weights of boys and girls (van Buuren et al., 2012) and median body weights of 18- to 79-year-old men and women were used, based on measured body heights of 16,500 men and 19,969 women in 13 EU Member States and assuming a BMI of 22 kg/m<sup>2</sup> (see Appendix 11 in EFSA NDA Panel (2013b)). The following growth factors have been applied: 0.25 for boys and girls aged 1–3 years, 0.06 for boys and girls aged 4–6 years, 0.13 for boys and girls aged 7–10 years, 0.11 for boys and 0.08 for girls aged 11–14 years and 0.08 for boys and 0.03 for girls aged 15–17 years. Growth factors were calculated as the proportional increase in protein requirement for growth relative to the maintenance requirement at the different ages (EFSA NDA Panel, 2012). The value for each age group corresponds to the mean of values for the years included (EFSA NDA Panel, 2014a). Calculated AIs were rounded to the nearest 10. Although the calculations yielded an AI for children aged 15–17 years that was higher (i.e. 410 mg/day) than the value set for adults (i.e. 400 mg/day), the Panel considered that there was no reason for such a difference, thus decided to set the same AI for children aged 15–17 years and adults.

The AIs for children are supported by total choline intake mean estimates in the EU (Section 3.2.1), i.e. estimates ranging from 151 to 210 mg/day (midpoint: 180 mg/day) in children aged 1 to < 3 years, from 177 to 304 mg/day (midpoint: 240 mg/day) in children aged 3 to < 10 years, from 244 to 373 mg/day (midpoint: 308 mg/day) among children aged 10 to < 18 years.

The Panel is aware that the AI for children aged 1–3 years (140 mg/day) is lower than the AI for infants aged 7–11 months (160 mg/day, Section 6.2). This difference is due to the approaches used for calculation (upwards extrapolation from the high choline intake of breast-fed infants from birth to 6 months, for infants aged 7–11 months vs downwards extrapolation from the AI for adults, for children aged 1–17 years). The Panel considers this higher AI for infants aged 7–11 months compared with children aged 1–3 years to be justified by a high demand for choline for phospholipid synthesis by the developing brain of infants (Section 2.3.4).

**Table 4:** Reference body weights and Adequate Intakes (AIs) of choline for children aged 1–17 years

Age (years)	Reference body weights (kg)		Growth factors		Calculated AIs (mg/day)		Calculated average AI (mg/day)	Proposed AIs (mg/day)
	Boys	Girls	Boys	Girls	Boys	Girls		
1–3	12.2 <sup>(a)</sup>	11.5 <sup>(a)</sup>	0.25	0.25	137.68	147.61	142.65	140
4–6	19.2 <sup>(b)</sup>	18.7 <sup>(b)</sup>	0.06	0.06	164.05	180.25	172.15	170
7–10	29.0 <sup>(c)</sup>	28.4 <sup>(c)</sup>	0.13	0.13	238.27	262.88	250.58	250
11–14	44.0 <sup>(d)</sup>	45.1 <sup>(d)</sup>	0.11	0.08	319.97	355.42	337.70	340
15–17	64.1 <sup>(e)</sup>	56.4 <sup>(e)</sup>	0.08	0.03	412.83	400.86	406.84	400 <sup>(f)</sup>

(a): Average of the median weight-for-age of male or female children aged 24 months according to the WHO Growth Standards (WHO Multicentre Growth Reference Study Group, 2006).

(b): Average of the median weight of male or female children aged 5 years (van Buuren et al., 2012).

(c): Average of the median weight of male or female children aged 8.5 years (van Buuren et al., 2012).

(d): Average of the median weight of male or female children aged 12.5 years (van Buuren et al., 2012).

(e): Average of the median weight of male or female children aged 16 years (van Buuren et al., 2012).

(f): The Panel decided to set the same AI for children aged 15–17 years and for adults.

Adult body weight used for calculations: 68.1 kg for men and 58.5 kg for women (median body weight of 18–79-year-old men and women, respectively, based on measured body heights of 16,500 men and 19,969 women in 13 EU Member States and assuming a BMI of 22 kg/m<sup>2</sup>, see Appendix 11 in EFSA NDA Panel (2013b)).

## 6.4. Pregnancy

The Panel concludes that calculation of choline transfer from the mother to the fetus and choline accretion in the fetus and placenta during the duration of pregnancy is not feasible to set DRVs for dietary choline during pregnancy due to a lack of data (Sections 2.3.3, 2.3.4 and 5.1.3.5). Although the available intervention studies on choline supplementation in the second half of pregnancy indicate that pregnant women may need more choline than non-pregnant women (Section 5.1.3.5), the data are not sufficient to allow an estimate of the additional requirement for dietary choline in pregnant women (above that of non-pregnant women).

Therefore, the Panel proposes to calculate the additional choline intake needed by pregnant woman, by isometric scaling from the AI of non-pregnant women (400 mg/day, Section 6.1), using the reference body weight for non-pregnant women and the mean gestational increase in body weight. The reference body weight of 18–79-year-old women (58.5 kg) was previously calculated from the measured body heights of 19,969 women in 13 EU Member States and assuming a BMI of 22 kg/m<sup>2</sup> (see Appendix 11 in EFSA NDA Panel (2013b)). A mean gestational increase in body weight of 12 kg, for women with a singleton pregnancy and a pre-pregnancy BMI in the range between 18.5 and 24.9 kg/m<sup>2</sup>, was also previously considered (EFSA NDA Panel, 2013b). Thus, the calculation was based on the equation below:

$$AI_{\text{pregnant}} = AI_{\text{non-pregnant}} \times \left( \frac{70.5 \text{ kg}}{58.5 \text{ kg}} \right) = 480 \text{ mg/day}$$

The Panel notes that the calculation by allometric scaling (as applied in Section 6.3) would lead to a value of 460 mg/day. The Panel, however, notes that the amount obtained by isometric scaling (480 mg/day) is the same as the lower dose in one intervention study on pregnant women (recruited at 27 weeks of gestation) (Yan et al., 2012) (Sections 2.3.6.1, 2.4.1.2 and 5.1.3). In view of the weak evidence and the minimal differences between the two scaling approaches, the Panel chose the value of 480 mg/day.

The Panel notes that this AI is higher than the mean choline intake of pregnant women (around 350 mg/day), observed either in the Latvian survey for which individual data were available to EFSA (Section 3.2.1) or in another publication outside the EU (Canada, Section 3.2.2).

The Panel proposes an AI of pregnant women of 480 mg choline/day. The Panel points out that this AI applies to the whole duration of pregnancy.



## 6.5. Lactation

The Panel concludes that the available intervention studies in lactating women (Sections 2.3.6.3 and 5.1.3.5) provide evidence that increased maternal choline intake enhances the concentration of choline in breast milk and that lactating women may need more choline than non-lactating women, but the data are not sufficient to allow an estimate of the additional requirement for dietary choline in lactating women (above that of non-lactating women).

For lactating women, the Panel decides to set a higher AI than for non-lactating women, by compensating for the secretion of choline in breast milk. Approximately 120 mg choline is secreted per day in human milk during the first 6 months of exclusive breastfeeding, considering an average concentration of total choline (free choline and choline compounds) in mature breast milk from mothers of full-term infants of 145 mg/L and a mean milk transfer during the first 6 months of lactation in exclusively breastfeeding women of 0.8 L/day (Section 2.3.6.3). The Panel proposes an additional AI of 120 mg/day above the AI for non-lactating women (400 mg/day), without correcting for intestinal absorption due to lack of data (Section 2.3.1). Thus, the Panel sets an AI of 520 mg/day for lactating women.

## Conclusions

The Panel considers that none of the biomarkers of choline intake or status is suitable to derive DRVs for choline. The Panel concludes that ARs and PRIs for choline cannot be derived for adults, infants and children, and therefore defines AIs. For all adults, the Panel sets an AI based on the midpoint of the range of observed mean choline intakes in healthy populations in the EU (about 370 mg/day), and in consideration of the results of a depletion/repletion study in which about 70% of the depleted subjects who had developed signs of organ dysfunction were repleted with an intake of about 400 mg/70-kg bw per day. For all infants aged 7–11 months, the Panel proposes an AI based on upwards extrapolation by allometric scaling from the estimated choline intake of exclusively breast-fed infants from birth to 6 months. For all children aged 1–17 years, the Panel derives AIs by downwards extrapolation from the adult AI, by allometric scaling, applying growth factors. These AIs are supported by estimated mean total choline intake in Europe. When applying allometric scaling, differences in reference body weight were taken into account. The Panel considers it unnecessary to give sex-specific AIs for adults, infants or children. For pregnant women, the Panel derives an AI by extrapolation from the AI for adults using isometric scaling and the mean gestational increase in body weight. For lactating women, the amount of choline secreted per day in human milk during the first 6 months of exclusive breastfeeding is added to the AI for non-lactating women (Table 5).

**Table 5:** Summary of Dietary Reference Values for choline

Age	Adequate Intakes (mg/day)
7–11 months	160
1–3 years	140
4–6 years	170
7–10 years	250
11–14 years	340
15–17 years	400
Adults	400
Pregnancy	480
Lactation	520

## Recommendations for research

The Panel suggests to undertake further research on:

- the identification of frequency of SNPs in genes coding for enzymes involved in choline metabolism that influence the requirement for dietary choline in the EU;
- the quantification of the increase in choline requirement, in carriers of alleles with increased need for choline;
- choline content of EU foods, to obtain better quantitative data on choline intake in Europe;

- biomarkers of choline status;
- criteria on which to base choline sufficiency in different populations;
- the consequences of the epigenetic modifications of genes involved in hormonal and vascular physiology and their expression following changes in choline intake during pregnancy;
- quantitative assessment of choline transfer from mother to fetus;
- quantification of the incorporated choline compounds in the body or in different organs during fetal development.

## References

- Abratte CM, Wang W, Li R, Moriarty DJ and Caudill MA, 2008. Folate intake and the MTHFR C677T genotype influence choline status in young Mexican American women. *Journal of Nutritional Biochemistry*, 19, 158–165.
- Abratte CM, Wang W, Li R, Axume J, Moriarty DJ and Caudill MA, 2009. Choline status is not a reliable indicator of moderate changes in dietary choline consumption in premenopausal women. *Journal of Nutritional Biochemistry*, 20, 62–69.
- Allen JC, Keller RP, Archer P and Neville MC, 1991. Studies in human lactation: milk composition and daily secretion rates of macronutrients in the first year of lactation. *American Journal of Clinical Nutrition*, 54, 69–80.
- Al-Waiz M, Mitchell SC, Idle JR and Smith RL, 1987. The metabolism of 14C-labelled trimethylamine and its N-oxide in man. *Xenobiotica*, 17, 551–558.
- Atkinson W, Elmslie J, Lever M, Chambers ST and George PM, 2008. Dietary and supplementary betaine: acute effects on plasma betaine and homocysteine concentrations under standard and postmethionine load conditions in healthy male subjects. *American Journal of Clinical Nutrition*, 87, 577–585.
- Au KS, Ashley-Koch A and Northrup H, 2010. Epidemiologic and genetic aspects of spina bifida and other neural tube defects. *Developmental Disabilities Research Reviews*, 16, 6–15.
- Bain MA, Fornasini G and Evans AM, 2005. Trimethylamine: metabolic, pharmacokinetic and safety aspects. *Current Drug Metabolism*, 6, 227–240.
- Baumgartner HK, Trinder KM, Galimanis CE, Post A, Phang T, Ross RG and Winn VD, 2015. Characterization of choline transporters in the human placenta over gestation. *Placenta*, 36, 1362–1369.
- Bayon Y, Croset M, Chirouze V, Tayot JL and Lagarde M, 1993. Phospholipid molecular species from human placenta lipids. *Lipids*, 28, 631–636.
- Bennett BJ, de Aguiar Vallim TQ, Wang Z, Shih DM, Meng Y, Gregory J, Allayee H, Lee R, Graham M, Crooke R, Edwards PA, Hazen SL and Lusis AJ, 2013. Trimethylamine-N-oxide, a metabolite associated with atherosclerosis, exhibits complex genetic and dietary regulation. *Cell Metabolism*, 17, 49–60.
- Bidulescu A, Chambless LE, Siega-Riz AM, Zeisel SH and Heiss G, 2007. Usual choline and betaine dietary intake and incident coronary heart disease: the Atherosclerosis Risk in Communities (ARIC) study. *BMC Cardiovascular Disorders*, 7, 20.
- Bitsanis D, Crawford MA, Moodley T, Holmsen H, Ghebremeskel K and Djahanbakhch O, 2005. Arachidonic acid predominates in the membrane phosphoglycerides of the early and term human placenta. *Journal of Nutrition*, 135, 2566–2571.
- Boeke CE, Gillman MW, Hughes MD, Rifas-Shiman SL, Villamor E and Oken E, 2013. Choline intake during pregnancy and child cognition at age 7 years. *American Journal of Epidemiology*, 177, 1338–1347.
- Boyd WD, Graham-White J, Blackwood G, Glen I and McQueen J, 1977. Clinical effects of choline in Alzheimer senile dementia. *Lancet*, 2, 711.
- Brody LC, Conley M, Cox C, Kirke PN, McKeever MP, Mills JL, Molloy AM, O'Leary VB, Parle-McDermott A, Scott JM and Swanson DA, 2002. A polymorphism, R653Q, in the trifunctional enzyme methylenetetrahydrofolate dehydrogenase/methenyltetrahydrofolate cyclohydrolase/formyltetrahydrofolate synthetase is a maternal genetic risk factor for neural tube defects: report of the Birth Defects Research Group. *American Journal of Human Genetics*, 71, 1207–1215.
- Buchman AL, Moukarzel A, Jenden DJ, Roch M, Rice K and Ament ME, 1993. Low plasma free choline is prevalent in patients receiving long term parenteral nutrition and is associated with hepatic aminotransferase abnormalities. *Clinical Nutrition*, 12, 33–37.
- Buchman AL, Dubin MD, Moukarzel AA, Jenden DJ, Roch M, Rice KM, Gornbein J and Ament ME, 1995. Choline deficiency: a cause of hepatic steatosis during parenteral nutrition that can be reversed with intravenous choline supplementation. *Hepatology*, 22, 1399–1403.
- Buchman AL, Sohel M, Moukarzel A, Bryant D, Schanler R, Awal M, Burns P, Dorman K, Belfort M, Jenden DJ, Killip D and Roch M, 2001. Plasma choline in normal newborns, infants, toddlers, and in very-low-birth-weight neonates requiring total parenteral nutrition. *Nutrition*, 17, 18–21.
- Butte NF, Garza C, Smith EO and Nichols BL, 1984. Human milk intake and growth in exclusively breast-fed infants. *Journal of Pediatrics*, 104, 187–195.
- Butte NF, Lopez-Alarcon MG and Garza C, 2002. *Nutrient adequacy of exclusive breastfeeding for the term infant during the first six months of life*. World Health Organization, Geneva, 47 pp.

- van Buuren S, Schönbeck Y and van Dommelen P, 2012. Collection, collation and analysis of data in relation to reference heights and reference weights for female and male children and adolescents (0–18 years) in the EU, as well as in relation to the age of onset of puberty and the age at which different stages of puberty are reached in adolescents in the EU. Project developed on the procurement project CT/EFSA/NDA/2010/01. EFSA supporting publication 2012:EN-255, 59 pp.
- Byrne CD and Targher G, 2014. Ectopic fat, insulin resistance, and nonalcoholic fatty liver disease: implications for cardiovascular disease. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 34, 1155–1161.
- Carmichael SL, Yang W and Shaw GM, 2010. Periconceptional nutrient intakes and risks of neural tube defects in California. *Birth Defects Research. Part A, Clinical and Molecular Teratology*, 88, 670–678.
- Caudill MA, Dellschaft N, Solis C, Hinkis S, Ivanov AA, Nash-Barboza S, Randall KE, Jackson B, Solomita GN and Vermeylen F, 2009. Choline intake, plasma riboflavin, and the phosphatidylethanolamine N-methyltransferase G5465A genotype predict plasma homocysteine in folate-deplete Mexican-American men with the methylenetetrahydrofolate reductase 677TT genotype. *Journal of Nutrition*, 139, 727–733.
- Chandra RK, 1984. Physical growth of exclusively breast-fed infants. *Nutrition Research*, 2, 275–276.
- Cheatham CL, Goldman BD, Fischer LM, da Costa KA, Reznick JS and Zeisel SH, 2012. Phosphatidylcholine supplementation in pregnant women consuming moderate-choline diets does not enhance infant cognitive function: a randomized, double-blind, placebo-controlled trial. *American Journal of Clinical Nutrition*, 96, 1465–1472.
- Chiuve SE, Giovannucci EL, Hankinson SE, Zeisel SH, Dougherty LW, Willett WC and Rimm EB, 2007. The association between betaine and choline intakes and the plasma concentrations of homocysteine in women. *American Journal of Clinical Nutrition*, 86, 1073–1081.
- Cho E, Zeisel SH, Jacques P, Selhub J, Dougherty L, Colditz GA and Willett WC, 2006. Dietary choline and betaine assessed by food-frequency questionnaire in relation to plasma total homocysteine concentration in the Framingham Offspring Study. *American Journal of Clinical Nutrition*, 83, 905–911.
- Cho E, Willett WC, Colditz GA, Fuchs CS, Wu K, Chan AT, Zeisel SH and Giovannucci EL, 2007a. Dietary choline and betaine and the risk of distal colorectal adenoma in women. *Journal of the National Cancer Institute*, 99, 1224–1231.
- Cho E, Holmes M, Hankinson SE and Willett WC, 2007b. Nutrients involved in one-carbon metabolism and risk of breast cancer among premenopausal women. *Cancer Epidemiology, Biomarkers and Prevention*, 16, 2787–2790.
- Cho E, Holmes MD, Hankinson SE and Willett WC, 2010. Choline and betaine intake and risk of breast cancer among post-menopausal women. *British Journal of Cancer*, 102, 489–494.
- Chu DM, Wahlqvist ML, Chang HY, Yeh NH and Lee MS, 2012. Choline and betaine food sources and intakes in Taiwanese. *Asia Pacific Journal of Clinical Nutrition*, 21, 547–557.
- Cohen BM, Renshaw PF, Stoll AL, Wurtman RJ, Yurgelun-Todd D and Babb SM, 1995. Decreased brain choline uptake in older adults. An *in vivo* proton magnetic resonance spectroscopy study. *JAMA*, 274, 902–907.
- Cole LK, Vance JE and Vance DE, 2012. Phosphatidylcholine biosynthesis and lipoprotein metabolism. *Biochimica et Biophysica Acta*, 1821, 754–761.
- Corbin KD and Zeisel SH, 2012. Choline metabolism provides novel insights into nonalcoholic fatty liver disease and its progression. *Current Opinion in Gastroenterology*, 28, 159–165.
- Corbin KD, Abdelmalek MF, Spencer MD, da Costa KA, Galanko JA, Sha W, Suzuki A, Guy CD, Cardona DM, Torquati A, Diehl AM and Zeisel SH, 2013. Genetic signatures in choline and 1-carbon metabolism are associated with the severity of hepatic steatosis. *FASEB Journal*, 27, 1674–1689.
- Cornford EM, Braun LD, Pardridge WM and Oldendorf WH, 1980. Blood flow rate and cellular influx of glucose and arginine in mouse liver *in vivo*. *American Journal of Physiology*, 238, H553–H560.
- da Costa KA, Gaffney CE, Fischer LM and Zeisel SH, 2005. Choline deficiency in mice and humans is associated with increased plasma homocysteine concentration after a methionine load. *American Journal of Clinical Nutrition*, 81, 440–444.
- da Costa KA, Kozyreva OG, Song J, Galanko JA, Fischer LM and Zeisel SH, 2006a. Common genetic polymorphisms affect the human requirement for the nutrient choline. *FASEB Journal*, 20, 1336–1344.
- da Costa KA, Niculescu MD, Craciunescu CN, Fischer LM and Zeisel SH, 2006b. Choline deficiency increases lymphocyte apoptosis and DNA damage in humans. *American Journal of Clinical Nutrition*, 84, 88–94.
- da Costa KA, Sanders LM, Fischer LM and Zeisel SH, 2011. Docosahexaenoic acid in plasma phosphatidylcholine may be a potential marker for *in vivo* phosphatidylethanolamine N-methyltransferase activity in humans. *American Journal of Clinical Nutrition*, 93, 968–974.
- da Costa KA, Corbin KD, Niculescu MD, Galanko JA and Zeisel SH, 2014. Identification of new genetic polymorphisms that alter the dietary requirement for choline and vary in their distribution across ethnic and racial groups. *FASEB Journal*, 28, 2970–2978.
- Craciun S and Balskus EP, 2012. Microbial conversion of choline to trimethylamine requires a glycol radical enzyme. *Proceedings of the National Academy of Sciences of the United States of America*, 109, 21307–21312.
- Cuddy LK, Winick-Ng W and Rylett RJ, 2014. Regulation of the high-affinity choline transporter activity and trafficking by its association with cholesterol-rich lipid rafts. *Journal of Neurochemistry*, 128, 725–740.

- Dalmeijer GW, Olthof MR, Verhoef P, Bots ML and van der Schouw YT, 2008. Prospective study on dietary intakes of folate, betaine, and choline and cardiovascular disease risk in women. *European Journal of Clinical Nutrition*, 62, 386–394.
- Davenport C and Caudill MA, 2013. Choline and milk. In: Zibadi S, Watson RR, Preedy VR (eds.). *Handbook of dietary and nutritional aspects of human breast milk. Human Health Handbooks volume 5*. Wageningen Academic Publishers, Wageningen, the Netherlands. pp. 335–352.
- Davenport C, Yan J, Taesuwan S, Shields K, West AA, Jiang X, Perry CA, Malysheva OV, Stabler SP, Allen RH and Caudill MA, 2015. Choline intakes exceeding recommendations during human lactation improve breast milk choline content by increasing PEMT pathway metabolites. *Journal of Nutritional Biochemistry*, 26, 903–911.
- Davies SEC, Woolf DA, Chalmers RA, Rafter JEM and Iles RA, 1992. Proton NMR studies of betaine excretion in the human neonate: consequences for choline and methyl group supply. *Journal of Nutritional Biochemistry*, 3, 523–530.
- De la Hueriga J and Popper H, 1951. Urinary excretion of choline metabolites following choline administration in normals and patients with hepatobiliary diseases. *Journal of Clinical Investigation*, 30, 463–470.
- DeLong CJ, Shen YJ, Thomas MJ and Cui Z, 1999. Molecular distinction of phosphatidylcholine synthesis between the CDP-choline pathway and phosphatidylethanolamine methylation pathway. *Journal of Biological Chemistry*, 274, 29683–29688.
- Dushianthan A, Goss V, Cusack R, Grocott M and Postle AD, 2014. Altered molecular specificity of surfactant phosphatidylcholine synthesis in patients with acute respiratory distress syndrome. *Respiratory Research*, 15, 128.
- EFSA (European Food Safety Authority), 2011. Use of the EFSA Comprehensive European Food Consumption Database in exposure assessment. *EFSA Journal* 2011;9(3):2097, 34 pp. doi:10.2903/j.efsa.2011.2097
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2009. Scientific Opinion on the appropriate age for introduction of complementary feeding of infants. *EFSA Journal* 2009;7(12):1423, 38 pp. doi: 10.2903/j.efsa.2009.1423
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2012. Scientific Opinion on Dietary Reference Values for protein. *EFSA Journal* 2012;10(2):2557, 66 pp. doi:10.2903/j.efsa.2012.2557
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2013a. Scientific Opinion on the safety of "citicoline" as a Novel Food ingredient. *EFSA Journal* 2013;11(10):3421, 22 pp. doi: 10.2903/j.efsa.2013.3421
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2013b. Scientific Opinion on Dietary Reference Values for energy. *EFSA Journal* 2013;11(1):3005, 112 pp. doi:10.2903/j.efsa.2013.3005
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2014a. Scientific Opinion on Dietary Reference Values for selenium. *EFSA Journal* 2014;12(10):3846, 66 pp. doi:10.2903/j.efsa.2014.3846
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2014b. Scientific Opinion on Dietary Reference Values for folate. *EFSA Journal* 2014;12(11):3893, 59 pp. doi:10.2903/j.efsa.2014.3893
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies), 2015. Scientific Opinion on Dietary Reference Values for cobalamin (vitamin B12). *EFSA Journal* 2015;13(7):4150, 64 pp. doi:10.2903/j.efsa.2015.4150
- Ehehalt R, Braun A, Karner M, Fullekrug J and Stremmel W, 2010. Phosphatidylcholine as a constituent in the colonic mucosal barrier—physiological and clinical relevance. *Biochimica et Biophysica Acta*, 1801, 983–993.
- El-Sohemy A, Xanthakos H, Beaulieu F, Allaire L and Fournier V, 2012. Literature search and review related to specific preparatory work in the establishment of Dietary References Values for thiamin, pantothenic acid and choline. Project developed on the procurement project CFT/EFSA/NUTRI/2011/01 (Lot 1). EFSA supporting publication. 229 pp.
- Enaw JO, Zhu H, Yang W, Lu W, Shaw GM, Lammer EJ and Finnell RH, 2006. CHKA and PCYT1A gene polymorphisms, choline intake and spina bifida risk in a California population. *BMC Medicine*, 4, 36.
- Fagone P and Jackowski S, 2013. Phosphatidylcholine and the CDP-choline cycle. *Biochimica et Biophysica Acta*, 1831, 523–532.
- FAO/WHO/UNU (Food and Agriculture Organization of the United Nations/World Health Organization/United Nations University), 2004. Human energy requirements. Report of a Joint FAO/WHO/UNU Expert Consultation: Rome, 17–24 October 2001. FAO Food and Nutrition Technical Report Series, 103 pp.
- Fayad LM, Salibi N, Wang X, Machado AJ, Jacobs MA, Bluemke DA and Barker PB, 2010. Quantification of muscle choline concentrations by proton MR spectroscopy at 3 T: technical feasibility. *AJR. American Journal of Roentgenology*, 194, W73–W79.
- Fischer LM, daCosta KA, Kwock L, Stewart PW, Lu TS, Stabler SP, Allen RH and Zeisel SH, 2007. Sex and menopausal status influence human dietary requirements for the nutrient choline. *American Journal of Clinical Nutrition*, 85, 1275–1285.
- Fischer LM, da Costa KA, Kwock L, Galanko J and Zeisel SH, 2010a. Dietary choline requirements of women: effects of estrogen and genetic variation. *American Journal of Clinical Nutrition*, 92, 1113–1119.
- Fischer LM, da Costa KA, Galanko J, Sha W, Stephenson B, Vick J and Zeisel SH, 2010b. Choline intake and genetic polymorphisms influence choline metabolite concentrations in human breast milk and plasma. *American Journal of Clinical Nutrition*, 92, 336–346.



- Gelenberg AJ, Doller-Wojcik JC and Growdon JH, 1979. Choline and lecithin in the treatment of tardive dyskinesia: preliminary results from a pilot study. *American Journal of Psychiatry*, 136, 772–776.
- Growdon JH, Cohen EL and Wurtman RJ, 1977. Huntington's disease: clinical and chemical effects of choline administration. *Annals of Neurology*, 1, 418–422.
- Hofvander Y, Hagman U, Hillervik C and Sjolín S, 1982. The amount of milk consumed by 1–3 months old breast- or bottle-fed infants. *Acta Paediatrica Scandinavica*, 71, 953–958.
- Holmes HC, Snodgrass GJ and Iles RA, 1996. The choline content of human breast milk expressed during the first few weeks of lactation. *Biochemical Society Transactions*, 24, 350S.
- Holmes HC, Snodgrass GJ and Iles RA, 2000. Changes in the choline content of human breast milk in the first 3 weeks after birth. *European Journal of Pediatrics*, 159, 198–204.
- Holmes-McNary MQ, Cheng WL, Mar MH, Fussell S and Zeisel SH, 1996. Choline and choline esters in human and rat milk and in infant formulas. *American Journal of Clinical Nutrition*, 64, 572–576.
- Ibiebele TI, Hughes MC, Pandeya N, Zhao Z, Montgomery G, Hayward N, Green AC, Whiteman DC, Webb PM, Study of Digestive H and Australian Cancer S, 2011. High intake of folate from food sources is associated with reduced risk of esophageal cancer in an Australian population. *Journal of Nutrition*, 141, 274–283.
- Icol YO, Ozbek R, Hamurtekin E and Ulus IH, 2005. Choline status in newborns, infants, children, breast-feeding women, breast-fed infants and human breast milk. *Journal of Nutritional Biochemistry*, 16, 489–499.
- IOM (Institute of Medicine), 1998. *Dietary Reference Intakes for thiamin, riboflavin, niacin, vitamin B6, folate, vitamin B12, pantothenic acid, biotin, and choline*. Food and Nutrition Board. National Academy Press, Washington, DC, USA, 591 pp.
- Ivanov A, Nash-Barboza S, Hinkis S and Caudill MA, 2009. Genetic variants in phosphatidylethanolamine N-methyltransferase and methylenetetrahydrofolate dehydrogenase influence biomarkers of choline metabolism when folate intake is restricted. *Journal of the American Dietetic Association*, 109, 313–318.
- Jacob RA, Jenden DJ, Allman-Farinelli MA and Swendseid ME, 1999. Folate nutriture alters choline status of women and men fed low choline diets. *Journal of Nutrition*, 129, 712–717.
- Jiang X, Yan J, West AA, Perry CA, Malysheva OV, Devapatla S, Pressman E, Vermeylen F and Caudill MA, 2012. Maternal choline intake alters the epigenetic state of fetal cortisol-regulating genes in humans. *FASEB Journal*, 26, 3563–3574.
- Jiang X, Bar HY, Yan J, Jones S, Brannon PM, West AA, Perry CA, Ganti A, Pressman E, Devapatla S, Vermeylen F, Wells MT and Caudill MA, 2013. A higher maternal choline intake among third-trimester pregnant women lowers placental and circulating concentrations of the antiangiogenic factor fms-like tyrosine kinase-1 (sFLT1). *FASEB Journal*, 27, 1245–1253.
- Johnson AR, Lao S, Wang T, Galanko JA and Zeisel SH, 2012. Choline dehydrogenase polymorphism rs12676 is a functional variation and is associated with changes in human sperm cell function. *PLoS ONE*, 7, e36047.
- Jope RS, Domino EF, Mathews BN, Sitaram N, Jenden DJ and Ortez A, 1982. Free and bound choline blood levels after phosphatidylcholine. *Clinical Pharmacology and Therapeutics*, 31, 483–487.
- Kim YI, Miller JW, da Costa KA, Nadeau M, Smith D, Selhub J, Zeisel SH and Mason JB, 1994. Severe folate deficiency causes secondary depletion of choline and phosphocholine in rat liver. *Journal of Nutrition*, 124, 2197–2203.
- Kohlmeier M, da Costa KA, Fischer LM and Zeisel SH, 2005. Genetic variation of folate-mediated one-carbon transfer pathway predicts susceptibility to choline deficiency in humans. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 16025–16030.
- Kotsopoulos J, Hankinson SE and Tworoger SS, 2010. Dietary betaine and choline intake are not associated with risk of epithelial ovarian cancer. *European Journal of Clinical Nutrition*, 64, 111–114.
- Lang DH, Yeung CK, Peter RM, Ibarra C, Gasser R, Itagaki K, Philpot RM and Rettie AE, 1998. Isoform specificity of trimethylamine N-oxygenation by human flavin-containing monooxygenase (FMO) and P450 enzymes: selective catalysis by FMO3. *Biochemical Pharmacology*, 56, 1005–1012.
- LASER Analytica, 2014. Comprehensive literature search and review of breast milk composition as preparatory work for the setting of dietary reference values for vitamins and minerals. Project developed on the procurement project RC/EFSA/NUTRI/2013/06 – OC/EFSA/SAS/2012/01. EFSA supporting publication 2014:EN-629, 154 pp.
- Lawrence CM, Millac P, Stout GS and Ward JW, 1980. The use of choline chloride in ataxic disorders. *Journal of Neurology, Neurosurgery and Psychiatry*, 43, 452–454.
- Lazo M, Hernaez R, Eberhardt MS, Bonekamp S, Kamel I, Guallar E, Koteish A, Brancati FL and Clark JM, 2013. Prevalence of nonalcoholic fatty liver disease in the United States: the Third National Health and Nutrition Examination Survey, 1988–1994. *American Journal of Epidemiology*, 178, 38–45.
- Lee JE, Giovannucci E, Fuchs CS, Willett WC, Zeisel SH and Cho E, 2010a. Choline and betaine intake and the risk of colorectal cancer in men. *Cancer Epidemiology, Biomarkers and Prevention*, 19, 884–887.
- Lee JE, Jacques PF, Dougherty L, Selhub J, Giovannucci E, Zeisel SH and Cho E, 2010b. Are dietary choline and betaine intakes determinants of total homocysteine concentration? *American Journal of Clinical Nutrition*, 91, 1303–1310.
- Lever M, Atkinson W, Sizeland PC, Chambers ST and George PM, 2007. Inter- and intra-individual variations in normal urinary glycine betaine excretion. *Clinical Biochemistry*, 40, 447–453.



- Lewis ED, Subhan FB, Bell RC, McCargar LJ, Curtis JM, Jacobs RL, Field CJ and team AP, 2014. Estimation of choline intake from 24 h dietary intake recalls and contribution of egg and milk consumption to intake among pregnant and lactating women in Alberta. *British Journal of Nutrition*, 112, 112–121.
- Li Z and Vance DE, 2008. Phosphatidylcholine and choline homeostasis. *Journal of Lipid Research*, 49, 1187–1194.
- Li Z, Agellon LB and Vance DE, 2007. Choline redistribution during adaptation to choline deprivation. *Journal of Biological Chemistry*, 282, 10283–10289.
- Lin CS and Wu RD, 1986. Choline oxidation and choline dehydrogenase. *Journal of Protein Chemistry*, 5, 193–200.
- Lockman PR and Allen DD, 2002. The transport of choline. *Drug Development and Industrial Pharmacy*, 28, 749–771.
- Lu MS, Fang YJ, Pan ZZ, Zhong X, Zheng MC, Chen YM and Zhang CX, 2015. Choline and betaine intake and colorectal cancer risk in Chinese population: a case-control study. *PLoS ONE*, 10, e0118661.
- Mazzetti S, Bracco C, Regge D, Caivano R, Russo F and Stasi M, 2013. Choline-containing compounds quantification by <sup>1</sup>H NMR spectroscopy using external reference and noise measurements. *Physica Medica*, 29, 677–683.
- Meck WH and Williams CL, 2003. Metabolic imprinting of choline by its availability during gestation: implications for memory and attentional processing across the lifespan. *Neuroscience and Biobehavioral Reviews*, 27, 385–399.
- Mehedint MG and Zeisel SH, 2013. Choline's role in maintaining liver function: new evidence for epigenetic mechanisms. *Current Opinion in Clinical Nutrition and Metabolic Care*, 16, 339–345.
- Michel V and Bakovic M, 2012. The ubiquitous choline transporter SLC44A1. *Central Nervous System Agents in Medicinal Chemistry*, 12, 70–81.
- Miller CA, Corbin KD, da Costa KA, Zhang S, Zhao X, Galanko JA, Blevins T, Bennett BJ, O'Connor A and Zeisel SH, 2014. Effect of egg ingestion on trimethylamine-N-oxide production in humans: a randomized, controlled, dose-response study. *American Journal of Clinical Nutrition*, 100, 778–786.
- Mischel W, 1956. Chemical composition of the human placenta with special consideration of the biogenous amine, choline. *Zentralblatt für Gynäkologie*, 78, 1089–1099.
- Mitchell SC and Smith RL, 2001. Trimethylaminuria: the fish malodor syndrome. *Drug Metabolism and Disposition*, 29, 517–521.
- Mygind VL, Evans SE, Peddie MC, Miller JC and Houghton LA, 2013. Estimation of usual intake and food sources of choline and betaine in New Zealand reproductive age women. *Asia Pacific Journal of Clinical Nutrition*, 22, 319–324.
- Neville MC, Keller R, Seacat J, Lutes V, Neifert M, Casey C, Allen J and Archer P, 1988. Studies in human lactation: milk volumes in lactating women during the onset of lactation and full lactation. *American Journal of Clinical Nutrition*, 48, 1375–1386.
- Niculescu MD, da Costa KA, Fischer LM and Zeisel SH, 2007. Lymphocyte gene expression in subjects fed a low-choline diet differs between those who develop organ dysfunction and those who do not. *American Journal of Clinical Nutrition*, 86, 230–239.
- O'Donoghue N, Sweeney T, Donagh R, Clarke KJ and Porter RK, 2009. Control of choline oxidation in rat kidney mitochondria. *Biochimica et Biophysica Acta*, 1787, 1135–1139.
- Okuda T and Haga T, 2000. Functional characterization of the human high-affinity choline transporter. *FEBS Letters*, 484, 92–97.
- Olthof MR, Brink EJ, Katan MB and Verhoef P, 2005. Choline supplemented as phosphatidylcholine decreases fasting and postmethionine-loading plasma homocysteine concentrations in healthy men. *American Journal of Clinical Nutrition*, 82, 111–117.
- Ouwerkerk R, Pettigrew RI and Gharib AM, 2012. Liver metabolite concentrations measured with <sup>1</sup>H MR spectroscopy. *Radiology*, 265, 565–575.
- Ozarda Ilcol Y, Uncu G and Ulus IH, 2002. Free and phospholipid-bound choline concentrations in serum during pregnancy, after delivery and in newborns. *Archives of Physiology and Biochemistry*, 110, 393–399.
- Ozarda Y, Cansev M and Ulus IH, 2014. Breast milk choline contents are associated with inflammatory status of breastfeeding women. *Breastfeeding Medicine*, 9, 39–41.
- Park EI and Garrow TA, 1999. Interaction between dietary methionine and methyl donor intake on rat liver betaine-homocysteine methyltransferase gene expression and organization of the human gene. *Journal of Biological Chemistry*, 274, 7816–7824.
- Pauwels S, Dopere I, Huybrechts I, Godderis L, Koppen G and Vansant G, 2015. Reproducibility and validity of an FFQ to assess usual intake of methyl-group donors. *Public Health Nutrition*, 18, 2530–2539.
- Poly C, Massaro JM, Seshadri S, Wolf PA, Cho E, Krall E, Jacques PF and Au R, 2011. The relation of dietary choline to cognitive performance and white-matter hyperintensity in the Framingham Offspring Cohort. *American Journal of Clinical Nutrition*, 94, 1584–1591.
- Pomfret EA, daCosta KA, Schurman LL and Zeisel SH, 1989. Measurement of choline and choline metabolite concentrations using high-pressure liquid chromatography and gas chromatography-mass spectrometry. *Analytical Biochemistry*, 180, 85–90.
- Pynn CJ, Henderson NG, Clark H, Koster G, Bernhard W and Postle AD, 2011. Specificity and rate of human and mouse liver and plasma phosphatidylcholine synthesis analyzed in vivo. *Journal of Lipid Research*, 52, 399–407.

- Reo NV, Adinezhadeh M and Foy BD, 2002. Kinetic analyses of liver phosphatidylcholine and phosphatidylethanolamine biosynthesis using  $(^{13}\text{C})$  NMR spectroscopy. *Biochimica et Biophysica Acta*, 1580, 171–188.
- Resseguie M, Song J, Niculescu MD, da Costa KA, Randall TA and Zeisel SH, 2007. Phosphatidylethanolamine N-methyltransferase (PEMT) gene expression is induced by estrogen in human and mouse primary hepatocytes. *FASEB Journal*, 21, 2622–2632.
- Resseguie ME, da Costa KA, Galanko JA, Patel M, Davis IJ and Zeisel SH, 2011. Aberrant estrogen regulation of PEMT results in choline deficiency-associated liver dysfunction. *Journal of Biological Chemistry*, 286, 1649–1658.
- Richman EL, Kenfield SA, Stampfer MJ, Giovannucci EL, Zeisel SH, Willett WC and Chan JM, 2012. Choline intake and risk of lethal prostate cancer: incidence and survival. *American Journal of Clinical Nutrition*, 96, 855–863.
- Rozen R, 1996. Molecular genetic aspects of hyperhomocysteinemia and its relation to folic acid. *Clinical and Investigative Medicine*, 19, 171–178.
- Savendahl L, Mar MH, Underwood LE and Zeisel SH, 1997. Prolonged fasting in humans results in diminished plasma choline concentrations but does not cause liver dysfunction. *American Journal of Clinical Nutrition*, 66, 622–625.
- SCF (Scientific Committee for Food), 1993. Nutrient and energy intakes for the European Community. Reports of the Scientific Committee for Food, 31st Series. Food - Science and Technique, European Commission, Luxembourg, 248 pp.
- Schwahn BC, Hafner D, Hohlfeld T, Balkenhol N, Laryea MD and Wendel U, 2003a. Pharmacokinetics of oral betaine in healthy subjects and patients with homocystinuria. *British Journal of Clinical Pharmacology*, 55, 6–13.
- Schwahn BC, Chen Z, Laryea MD, Wendel U, Lussier-Cacan S, Genest J Jr, Mar MH, Zeisel SH, Castro C, Garrow T and Rozen R, 2003b. Homocysteine-betaine interactions in a murine model of 5,10-methylenetetrahydrofolate reductase deficiency. *FASEB Journal*, 17, 512–514.
- Sha W, da Costa KA, Fischer LM, Milburn MV, Lawton KA, Berger A, Jia W and Zeisel SH, 2010. Metabolomic profiling can predict which humans will develop liver dysfunction when deprived of dietary choline. *FASEB Journal*, 24, 2962–2975.
- Shaw GM, Carmichael SL, Yang W, Selvin S and Schaffer DM, 2004. Periconceptional dietary intake of choline and betaine and neural tube defects in offspring. *American Journal of Epidemiology*, 160, 102–109.
- Solis C, Veenema K, Ivanov AA, Tran S, Li R, Wang W, Moriarty DJ, Maletz CV and Caudill MA, 2008. Folate intake at RDA levels is inadequate for Mexican American men with the methylenetetrahydrofolate reductase 677TT genotype. *Journal of Nutrition*, 138, 67–72.
- Song J, da Costa KA, Fischer LM, Kohlmeier M, Kwock L, Wang S and Zeisel SH, 2005. Polymorphism of the PEMT gene and susceptibility to nonalcoholic fatty liver disease (NAFLD). *FASEB Journal*, 19, 1266–1271.
- Spencer MD, Hamp TJ, Reid RW, Fischer LM, Zeisel SH and Fodor AA, 2011. Association between composition of the human gastrointestinal microbiome and development of fatty liver with choline deficiency. *Gastroenterology*, 140, 976–986.
- Stead LM, Brosnan JT, Brosnan ME, Vance DE and Jacobs RL, 2006. Is it time to reevaluate methyl balance in humans? *American Journal of Clinical Nutrition*, 83, 5–10.
- Svennerholm L and Vanier MT, 1972. The distribution of lipids in the human nervous system. II. Lipid composition of human fetal and infant brain. *Brain Research*, 47, 457–468.
- Sweiry JH, Page KR, Dacke CG, Abramovich DR and Yudilevich DL, 1986. Evidence of saturable uptake mechanisms at maternal and fetal sides of the perfused human placenta by rapid paired-tracer dilution: studies with calcium and choline. *Journal of Developmental Physiology*, 8, 435–445.
- Tang WH, Wang Z, Levison BS, Koeth RA, Britt EB, Fu X, Wu Y and Hazen SL, 2013. Intestinal microbial metabolism of phosphatidylcholine and cardiovascular risk. *New England Journal of Medicine*, 368, 1575–1584.
- Ueland PM, 2011. Choline and betaine in health and disease. *Journal of Inherited Metabolic Disease*, 34, 3–15.
- Ueland PM, Holm PI and Hustad S, 2005. Betaine: a key modulator of one-carbon metabolism and homocysteine status. *Clinical Chemistry and Laboratory Medicine*, 43, 1069–1075.
- USDA (US Department of Agriculture), 2012. Nutrient intakes from food: mean amounts consumed per individual, by gender and age, what we eat in America, NHANES 2009–2010. Agricultural Research Service. Available online: [http://www.ars.usda.gov/SP2UserFiles/Place/12355000/pdf/0910/Table\\_1\\_NIN\\_GEN\\_09.pdf](http://www.ars.usda.gov/SP2UserFiles/Place/12355000/pdf/0910/Table_1_NIN_GEN_09.pdf)
- USDA (US Department of Agriculture), 2013. USDA National Nutrient Database for Standard Reference, release 26. Agricultural Research Service. Available online: <http://ndb.nal.usda.gov/ndb/nutrients/report/nutrientsfrm?max=25&offset=0&totCount=0&nutrient1=421 &nutrient2=&nutrient3=&subset=0&fg= &sort=f&measureby=g>
- USDA (US Department of Agriculture), 2015. USDA National Nutrient Database for Standard Reference, release 28. Agricultural Research Service. Available online: <http://ndb.nal.usda.gov/ndb/nutrients/report/nutrientsfrm?max=25&offset=0&totCount=0 &nutrient1=421&nutrient2= &nutrient3=&subset=0&fg=&sort=f &measureby=g>
- Vance DE, 2014. Phospholipid methylation in mammals: from biochemistry to physiological function. *Biochimica et Biophysica Acta*, 1838, 1477–1487.
- Vance DE, Li ZY and Jacobs RL, 2007. Hepatic phosphatidylethanolamine N-methyltransferase, unexpected roles in animal biochemistry and physiology. *Journal of Biological Chemistry*, 282, 33237–33241.
- Varela-Moreiras G, Selhub J, da Costa KA and Zeisel SH, 1992. Effect of chronic choline deficiency in rats on liver folate content and distribution. *The Journal of Nutritional Biochemistry*, 3, 519–522.

- Veenema K, Solis C, Li R, Wang W, Maletz CV, Abratte CM and Caudill MA, 2008. Adequate Intake levels of choline are sufficient for preventing elevations in serum markers of liver dysfunction in Mexican American men but are not optimal for minimizing plasma total homocysteine increases after a methionine load. *American Journal of Clinical Nutrition*, 88, 685–692.
- Vennemann FB, Ioannidou S, Valsta LM, Dumas C, Ocke MC, Mensink GB, Lindtner O, Virtanen SM, Tlustos C, D'Addezio L, Mattison I, Dubuisson C, Siksnia I and Heraud F, 2015. Dietary intake and food sources of choline in European populations. *British Journal of Nutrition*, 114, 2046–2055.
- Villamor E, Rifas-Shiman SL, Gillman MW and Oken E, 2012. Maternal intake of methyl-donor nutrients and child cognition at 3 years of age. *Paediatric and Perinatal Epidemiology*, 26, 328–335.
- Visentin CE, Masih S, Plumptre L, Malysheva O, Nielsen DE, Sohn KJ, Ly A, Lausman AY, Berger H, Croxford R, El-Sohehy A, Caudill MA, O'Connor DL and Kim YI, 2015. Maternal Choline Status, but Not Fetal Genotype, Influences Cord Plasma Choline Metabolite Concentrations. *Journal of Nutrition*, 145, 1491–1497.
- Wallace JM, McCormack JM, McNulty H, Walsh PM, Robson PJ, Bonham MP, Duffy ME, Ward M, Molloy AM, Scott JM, Ueland PM and Strain JJ, 2012. Choline supplementation and measures of choline and betaine status: a randomised, controlled trial in postmenopausal women. *British Journal of Nutrition*, 108, 1264–1271.
- Wang Z, Klipfell E, Bennett BJ, Koeth R, Levison BS, Dugar B, Feldstein AE, Britt EB, Fu X, Chung YM, Wu Y, Schauer P, Smith JD, Allayee H, Tang WH, DiDonato JA, Lusis AJ and Hazen SL, 2011. Gut flora metabolism of phosphatidylcholine promotes cardiovascular disease. *Nature*, 472, 57–63.
- Wang Z, Tang WH, Buffa JA, Fu X, Britt EB, Koeth RA, Levison BS, Fan Y, Wu Y and Hazen SL, 2014. Prognostic value of choline and betaine depends on intestinal microbiota-generated metabolite trimethylamine-N-oxide. *European Heart Journal*, 35, 904–910.
- Welsch F, 1976. Studies on accumulation and metabolic fate of (N-Me<sub>3</sub>)choline in human term placenta fragments. *Biochemical Pharmacology*, 25, 1021–1030.
- West AA, Yan J, Jiang X, Perry CA, Innis SM and Caudill MA, 2013. Choline intake influences phosphatidylcholine DHA enrichment in nonpregnant women but not in pregnant women in the third trimester. *American Journal of Clinical Nutrition*, 97, 718–727.
- WHO Multicentre Growth Reference Study Group (World Health Organization), 2006. WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development. 312 pp.
- Widdowson EM, 1963. Growth and composition of the fetus and newborn. In: Assali NS (ed). *Biology of gestation*. Academic Press, New York, NY. pp. 1–51.
- Wilcken DE, Wang XL, Sim AS and McCredie RM, 1996. Distribution in healthy and coronary populations of the methylenetetrahydrofolate reductase (MTHFR) C677T mutation. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 16, 878–882.
- Wu BT, Dyer RA, King DJ, Richardson KJ and Innis SM, 2012. Early second trimester maternal plasma choline and betaine are related to measures of early cognitive development in term infants. *PLoS ONE*, 7, e43448.
- Xu X, Gammon MD, Zeisel SH, Lee YL, Wetmur JG, Teitelbaum SL, Bradshaw PT, Neugut AI, Santella RM and Chen J, 2008. Choline metabolism and risk of breast cancer in a population-based study. *FASEB Journal*, 22, 2045–2052.
- Xu X, Gammon MD, Zeisel SH, Bradshaw PT, Wetmur JG, Teitelbaum SL, Neugut AI, Santella RM and Chen J, 2009. High intakes of choline and betaine reduce breast cancer mortality in a population-based study. *FASEB Journal*, 23, 4022–4028.
- Yan J, Wang W, Gregory JF 3rd, Malysheva O, Brenna JT, Stabler SP, Allen RH and Caudill MA, 2011. MTHFR C677T genotype influences the isotopic enrichment of one-carbon metabolites in folate-compromised men consuming d<sub>9</sub>-choline. *American Journal of Clinical Nutrition*, 93, 348–355.
- Yan J, Jiang X, West AA, Perry CA, Malysheva OV, Devapatla S, Pressman E, Vermeylen F, Stabler SP, Allen RH and Caudill MA, 2012. Maternal choline intake modulates maternal and fetal biomarkers of choline metabolism in humans. *American Journal of Clinical Nutrition*, 95, 1060–1071.
- Yan J, Jiang X, West AA, Perry CA, Malysheva OV, Brenna JT, Stabler SP, Allen RH, Gregory JF 3rd and Caudill MA, 2013. Pregnancy alters choline dynamics: results of a randomized trial using stable isotope methodology in pregnant and nonpregnant women. *American Journal of Clinical Nutrition*, 98, 1459–1467.
- Yu D, Shu XO, Xiang YB, Li H, Yang G, Gao YT, Zheng W and Zhang X, 2014. Higher dietary choline intake is associated with lower risk of nonalcoholic fatty liver in normal-weight Chinese women. *Journal of Nutrition*, 144, 2034–2040.
- Zeisel SH, 2006. Choline: critical role during fetal development and dietary requirements in adults. *Annual Review of Nutrition*, 26, 229–250.
- Zeisel SH, 2007. Gene response elements, genetic polymorphisms and epigenetics influence the human dietary requirement for choline. *IUBMB Life*, 59, 380–387.
- Zeisel SH, 2012. Dietary choline deficiency causes DNA strand breaks and alters epigenetic marks on DNA and histones. *Mutation Research*, 733, 34–38.
- Zeisel SH and Blusztajn JK, 1994. Choline and human nutrition. *Annual Review of Nutrition*, 14, 269–296.
- Zeisel SH, Growdon JH, Wurtman RJ, Magil SG and Logue M, 1980. Normal plasma choline responses to ingested lecithin. *Neurology*, 30, 1226–1229.

- Zeisel SH, Char D and Sheard NF, 1986. Choline, phosphatidylcholine and sphingomyelin in human and bovine milk and infant formulas. *Journal of Nutrition*, 116, 50–58.
- Zeisel SH, Da Costa KA, Franklin PD, Alexander EA, Lamont JT, Sheard NF and Beiser A, 1991. Choline, an essential nutrient for humans. *FASEB Journal*, 5, 2093–2098.
- Zeisel SH, Mar MH, Howe JC and Holden JM, 2003. Concentrations of choline-containing compounds and betaine in common foods. *Journal of Nutrition*, 133, 1302–1307.
- Zhang AQ, Mitchell SC and Smith RL, 1999. Dietary precursors of trimethylamine in man: a pilot study. *Food and Chemical Toxicology*, 37, 515–520.

## Abbreviations

AI	Adequate Intake
ALT	alanine aminotransferase
AR	Average Requirement
ARA	arachidonic acid
AST	aspartate aminotransferase
ATP	adenosine triphosphate
BADH	betaine aldehyde dehydrogenase
BE	Barrett's oesophagus
BHMT	betaine-homocysteine methyltransferase
BMI	body mass index
bw	body weight
CCT	phosphocholine cytidyltransferase
CDP	cytidine 5-diphosphate
CEPT	choline/ethanolaminephosphotransferase
CHK	choline kinase
CHKA	choline kinase A
CHKB	choline kinase B
CHD	coronary heart disease
CHDH	choline dehydrogenase or choline oxidase
CHT	choline transporter
CI	confidence interval
CK	creatine (phospho)kinase
COMET	single-cell gel electrophoresis
CpG	cytosine–phosphate–guanine
CPT	cytidine 5-diphosphate-choline
CRC	colorectal cancer
CRH	corticotropin releasing hormone
CRP	C-reactive protein
CT	computerised tomography
CTL1	choline transporter-like protein 1
CTP	cytidine triphosphate
CVD	cardiovascular disease
DFE	dietary folate equivalent
DHA	docosahexaenoic acid
DMG	dimethylglycine
DNA	Deoxyribonucleic acid
DRV	Dietary Reference Value
dTMP	deoxythymidine monophosphate
EAR	Estimated Average Requirement
FFQ	food frequency questionnaire
FMO3	flavin-containing monooxygenase isoform 3
GC–MS	gas chromatography–mass spectrometry
GGT	γ-glutamyltransferase
GPC	glycerophosphocholine
HClO <sub>4</sub>	perchloric acid
Hcy	homocysteine
HDL	high-density lipoprotein
HPLC	high-performance liquid chromatography

HR	hazard ratio
IOM	US Institute of Medicine of the National Academy of Sciences
KBIT-2	Kaufmann Brief Intelligence Test, Second Edition
$K_m$	Michaelis constant
LC-MS/MS	liquid chromatography–tandem mass spectrometry
LC-PUFA	long-chain polyunsaturated fatty acid
LDH	lactate dehydrogenase
LDL	low-density lipoprotein
LOAEL	Lowest Observed Adverse Effect Level
Met	methionine
MG	methylglycine
MIDA	multiple isotopomer distribution analysis
MM	molecular mass
MRI	magnetic resonance imaging
mRNA	messenger RNA
MRS	magnetic resonance spectroscopy
MS	methionone synthase
MTHFD1	5,10-methylenetetrahydrofolate dehydrogenase 1
MTHFR	methylenetetrahydrofolate reductase
NAFLD	non-alcoholic fatty liver disease
NDA	EFSA Panel on Dietetic Products, Nutrition and Allergies
NHS	Nurses' Health Study
NHANES	National Health and Nutrition Examination Survey
NORCCAP	Norwegian Colorectal Cancer Prevention
NR3C1	nuclear receptor subfamily 3 group C member 1 (glucocorticoid receptor)
N.S.	not significant
NTD	neural tube defect
OCT	organic cation transporter
OR	odds ratio
P	percentile
PC	phosphatidylcholine
PChol	phosphocholine
PE	phosphatidylethanolamine
PEMT	phosphatidylethanolamine <i>N</i> -methyltransferase
PL	phospholipase
PRI	Population Reference Intake
Q	quintile
RCT	randomised controlled trial
RDA	Recommended Dietary Allowance
RFC1	reduced folate carrier 1
RNA	ribonucleic acid
RR	relative risk
rs number	Reference SNP cluster ID
SAH	<i>S</i> -adenosylhomocysteine
SAH-H	<i>S</i> -adenosylhomocysteine hydrolase
SAM	<i>S</i> -adenosyl-methionine
SCF	Scientific Committee for Food
SD	standard deviation
SEM	standard error of the mean
sFLT1	soluble fms-like tyrosine kinase 1
SLC44A1	solute carrier family 44 member 1 (choline transporter)
SLC5A7	solute carrier family 5 member 7 (choline transporter)
SNP	single nucleotide polymorphism
SPM	sphingomyelin
TAG	triacylglycerol
tHcy	total homocysteine
THF	tetrahydrofolate



TLC	thin-layer chromatography
TMA	trimethylamine
TMAO	trimethylamine- <i>N</i> -oxide
TMP	thymidine monophosphate (thymidylic acid)
TTMA	total trimethylamine
TUNEL	terminal deoxynucleotidyl transferase dUTP nick end labeling
UL	Tolerable Upper Intake Level
USDA	United States Department of Agriculture
VLDL	very low-density lipoproteins
WHO	World Health Organization
WRAML2	Wide Range Assessment of Memory and Learning Second Edition

## Appendix A – Concentrations of free and total choline in breast milk of healthy lactating mothers

Reference	Number of women (number of samples)	Country	Maternal dietary intake (mg/day) Mean $\pm$ SE (range)	Stage of lactation	Choline concentration (mg/L)		Analytical method	Comments
					Mean $\pm$ SE	Median		
Holmes-McNary et al. (1996)	16(16)	USA	Not reported	27–32 days post partum	Free choline $12.1 \pm 2.3$ Total choline <sup>(a)</sup> $130.6 \pm 25.3$		Water-soluble compounds extracted with perchloric acid (HClO <sub>4</sub> ), HPLC after hydrolysis; phospholipid-bound choline separated by TLC and analysed after hydrolysis by GC-MS or phosphorus quantification	Hospital bank milk Pumped milk samples Full-term infants No information on polymorphism and supplementation of the mothers Plasma choline concentration was not assessed
Holmes et al. (2000)	8(8)	UK	Not reported	2–6 days post partum 7–22 days post partum	Free choline $11 \pm 2$ Total choline <sup>(a)</sup> $63 \pm 9$ Free choline $22 \pm 5$ Total choline <sup>(a)</sup> $133 \pm 15$		Nuclear magnetic resonance spectrometry (extraction with HClO <sub>4</sub> and chloroform of water soluble and phospholipid-bound choline, respectively)	Infants born at 28–38 weeks of gestation (preterm and term) No information on the supplementation of the mothers Aliquots of expressed foremilk Plasma choline concentration not reported
İlcol et al. (2005)	(21)	Turkey	Not reported	Colostrum (0–2 days after birth)	Free choline $13.8 \pm 2.2$ Total choline <sup>(a)</sup> $70.4 \pm 3.6$		* <b>Free choline</b> in milk: measured with a modification of the enzymatic radiochemical method	116 breastfeeding women : 32 smokers No information about the term of the infants and

Reference	Number of women (number of samples)	Country	Maternal dietary intake (mg/day) Mean $\pm$ SE (range)	Stage of lactation	Choline concentration (mg/L)		Analytical method	Comments
					Mean $\pm$ SE	Median		
	(95)			12–180 days post partum	Free choline $23.8 \pm 1.04$ Total choline <sup>(a)</sup> $153.8 \pm 5.0$		*Phospholipid-bound choline, PC and SPM in milk: measured with an enzymatic colorimetric method *PChol and GPC: first hydrolysed enzymatically to free choline then measured with high-performance liquid chromatography–electrochemical detection system	supplementation of the mothers For colostrum analyses 0–2 days, 57 full-term plus 24 preterm infants were investigated; milks from day 12–180 were provided by 95 mothers with no indication of gestational age Maternal plasma choline concentration reported and correlation with breast milk concentration investigated Inverse linear relationship between free choline concentration in breast milk and lactating days of the mothers ( $r = -0.625$ ; $p < 0.001$ )
	(14)		12–28 days post partum	Free choline $31.1 \pm 3.8$ Total choline <sup>(a)</sup> $166.2 \pm 8.5$				
	(12)		75–90 days post partum	Free choline $29.8 \pm 2.2$ Total choline <sup>(a)</sup> $150.1 \pm 8.8$				
	(11)			165–180 days post partum	Free choline $13.8 \pm 1.6$ Total choline <sup>(a)</sup> $140.5 \pm 10.9$ All breast milk free choline and total choline mean values were significantly higher than colostrum values, except free choline value for days 165–180 which was significantly lower than the value for days 12–180			

Reference	Number of women (number of samples)	Country	Maternal dietary intake (mg/day) Mean ± SE (range)	Stage of lactation	Choline concentration (mg/L)		Analytical method	Comments
					Mean ± SE	Median		
Fischer et al. (2010b)	51(51)	USA American (89%), African-American (3%), Asian (6%), American Indian (1%), other (1%)	<u>Supplemented group (n = 48)</u> (Supplement: 750 mg choline/day) *Dietary choline: 338 ± 14 (124–622) *Total choline intake: 1,088 ± 14	45 days post partum	Free choline 11.0 ± 1.0 (Significantly higher than in placebo group) <u>Total choline<sup>(a)</sup></u> 149.4		Liquid chromatography/electrospray ionisation isotope dilution mass spectrometry	103 participants: no breast milk data for four individuals and no dietary intakes for nine individuals 3 days dietary records at 45 days post partum PC supplement or placebo from 18 weeks of gestation to 90 days post partum Calculated duration of pregnancy (from duration of treatment) 34–42 weeks (for supplementation group) and 35–43 weeks (for placebo group) Maternal plasma choline concentration reported Genetic polymorphism investigated Correlation between breast milk concentration of choline or plasma concentration of choline and total choline intake
	48(48)		<u>Placebo group (n = 46)</u> * Dietary choline: 364 ± 18 (139–671) *Total choline intake: 364 ± 18	45 days post partum	Free choline 8.6 ± 0.8 <u>Total choline<sup>(a)</sup></u> 124.8			

Reference	Number of women (number of samples)	Country	Maternal dietary intake (mg/day) Mean $\pm$ SE (range)	Stage of lactation	Choline concentration (mg/L)		Analytical method	Comments
					Mean $\pm$ SE	Median		
Ozarda et al. (2014)	53	Turkey	Not reported	1–3 days post partum	Free choline 7.4 (2.2–13.6) <sup>(c)</sup> Total choline 42.4 (31.4–72.1) <sup>(b),(c)</sup>		HPLC–electrochemical detection for water-soluble choline compounds after hydrolysis; phospholipid-bound choline by enzymatic colorimetric method	Women who provided colostrum samples were not the same as the women who provided the mature milk samples Term infants Expressed milk Supplementation of the mothers not reported Plasma CRP concentration reported (relationship between serum CRP and breast milk content investigated)
	54			22–180 days post partum	Free choline 9.7 (7.0–13.9) <sup>(c)</sup> Total choline 159.6 (130.2–176.6) <sup>(b),(c)</sup> Free and total choline median values at days 1–3 were significantly lower than at days 22–180. In colostrum, positive correlation of water-soluble choline compounds with CRP in maternal serum and negative correlation with PC. No such correlation in mature milk			



Reference	Number of women (number of samples)	Country	Maternal dietary intake (mg/day) Mean ± SE (range)	Stage of lactation	Choline concentration (mg/L)		Analytical method	Comments
					Mean ± SE	Median		
Davenport et al. (2015)	28	USA	*Dietary choline: 380 <b>(a) Supplement: 100</b> *Total choline intake: 480 (n = 15) <b>(b) Supplement: 550</b> *Total choline intake: 930 (n = 13)	5 weeks post partum	Mean ± SD  (a) <u>Free choline</u> Baseline: 8.9 ± 4.2 Week 10: 16.5 ± 1.3 (a) <u>Total choline<sup>(a)</sup></u> Baseline: 136.5 ± 26.0 Week 10: 104.2 ± 5.2 (b) <u>Free choline</u> Baseline: 8.8 ± 5.7 Week 10: 15.4 ± 1.4 (b) <u>Total choline<sup>(a)</sup></u> Baseline: 117.1 ± 22.8 Week 10: 125.0 ± 6.3 <b>All subjects</b> <u>Free choline:</u> Baseline: 8.8 ± 4.4 <u>Total choline<sup>(a)</sup></u> Baseline: 127.5 ± 26.0	LC-MS/MS	No information about the term of the infants Expressed milk Maternal plasma choline concentration reported Correlation between breast milk concentration of choline and total choline intake Increased circulating plasma choline during lactation The study also had a control group (non-pregnant, non-lactating women)	

CRP: C-reactive protein; GC-MS: gas chromatography-mass spectrometry; GPC: glycerophosphocholine; HClO4: perchloric acid; HPLC: high-performance liquid chromatography; LC-MS/MS: liquid chromatography-tandem mass spectrometry; PC: phosphatidylcholine; PChol: phosphocholine; SE: standard error; SPM: sphingomyelin; TLC: thin-layer chromatography.

(a): Total choline was the result of the sum of: free choline, phosphatidylcholine, phosphocholine, glycerophosphocholine, sphingomyelin.

(b): Total choline was the result of the sum of: free choline, phosphocholine, glycerophosphocholine, phospholipid-bound choline.

(c): Median (P25-P75).

The values of free choline and total choline concentration in breast milk reported in the articles were expressed in nmol/mL or mmol/L, those values were converted in mg/L using the following molecular mass (MW) (for free choline and total choline) = 104.17 g/mol.

## Appendix B – Intervention and observational studies on the relationship between dietary choline and plasma homocysteine concentration

Author	Type of study	Subjects n, sex, age, country	*Intervention/ design (trials) *Intake measurement (cross-sectional studies)	*Duration (trials) *Choline intake (mg/day) (cross-sectional studies)	tHcy in plasma (µmol/L)		Comment on tHcy in plasma	Other outcomes
					Plasma tHcy (fasting)	Post-methionine (0.1 g/kg bw)		
Olthof et al. (2005)	Double-blind cross-over RCT	26 (men), 50-71 years, NL	<b>2.6 g choline/day</b> as PC, n = 13	2 weeks	Mean ± SD Baseline 15.6 ± 4.0 Day 15 13.6 ± 2.5	Mean ± SD Baseline 27.0 ± 6.1 Day 15 22.3 ± 3.3	Choline 2.6 g/ day for 2 weeks decreased significantly fasting plasma tHcy and 6-h post-methionine plasma tHcy	Choline supplement decreased serum folate and alkaline phosphatase, increased serum B6 and TAG; no change in cobalamin, ALT, AST, GGT, creatinine, total, LDL and HDL cholesterol
			No supplement (wash-out period) Placebo, n = 13	2 weeks	–	–		
Wallace et al. (2012)	Double-blind RCT	42 (postmenopausal women), 49-71 years, Ireland	<b>1 g choline/day</b> (as bitartrate), n = 19	12 weeks	Median Baseline 9.9 6 weeks 9.5 12 weeks 9.7		No significant difference of plasma tHcy at 6 and 12 weeks	MTHFR genotype TT 10.5% in choline group Plasma choline, betaine and DMG at 6 weeks significantly higher in choline group than placebo group MTHFR genotype TT 4.3% in placebo group
			Placebo (2.4 g tartaric acid)/day, n = 23		Baseline 9.7 6 weeks 10.1 12 weeks 10.0			
Atkinson et al. (2008)	Randomised, single-event, cross-over	8 (men), 19-40 years, New Zealand	<b>500 mg choline</b> as chloride	Once per week	Non-significant decrease		Choline from a meal has a greater tHcy lowering effect than supplemental	Increase in plasma betaine except after the low-choline/low- betaine meal Urinary betaine excretion did not
			High-choline meal <b>(760 mg choline)</b>	Once per week	Significant decrease by 0.77 µmol after 4-6 h			

Author	Type of study	Subjects n, sex, age, country	*Intervention/ design (trials) *Intake measurement (cross-sectional studies)	*Duration (trials) *Choline intake (mg/day) (cross-sectional studies)	tHcy in plasma (µmol/L)		Comment on tHcy in plasma	Other outcomes
					Plasma tHcy (fasting)	Post-methionine (0.1 g/kg bw)		
			High-choline meal <b>(760 mg choline)</b> plus methionine load (100 mg/kg bw)	Once per week		Significant lower rise at 4–6 h compared to low- choline meal by 6.9–7.6 µmol/L	change Urinary DMG excretion increased after high-choline meal, not after choline supplement	
			Low-choline meal <b>(&lt; 1 mg choline)</b>	Once per week				
Cho et al. (2006)	Cross- sectional study in a long-term cohort, offspring of Framingham cohort, start 1971, 5th examination 1991–1994	1,860, (1,040 women), 28- 82 years, USA	FFQ	Energy-adjusted intake Total choline (all forms): <b>313</b> ± 61 (mean ± SD)	Adjusted for age, sex, folate, B6, cobalamin intake, smoking, alcohol, caffeine, medication, serum creatinine	Hcy lowering effect observed at choline intakes < 1,000 mg/day and stronger in men than in women		
				Quintiles (mean)	Geometric mean (95% CI)			
				Q1 <b>234</b>	10.6 (10.2, 11.0)			
				Q2 <b>283</b>	10.4 (10.0, 10.8)			
				Q3 <b>311</b>	10.1 (9.7, 10.5)			
				Q4 <b>339</b>	9.7 (9.3, 10.1)			
				Q5 <b>401</b>	9.8 (9.5, 10.2)			
Lee et al. (2010b) (follow-up from Cho et al. (2006))	Cross- sectional study in long- term cohort study, Framingham Offspring study started 1971-1974	2,732 (1,325 men), 29–86 years, USA	FFQ, 6th examination 1995–1998	Energy-adjusted total intake Total choline (all forms): <b>308</b> ± 56 (mean ± SD)	Adjusted for age, sex, folate, B6, cobalamin intake, smoking, alcohol, caffeine, total energy, serum creatinine	Inverse association between choline intake and either fasting or post- methionine plasma tHcy before folic acid fortification in the USA, not after Association	Choline intake quintiles differ from 1 to 5 by 145 mg/day only	
				Quintiles (median ± SD)	Geometric mean (95% CI)			
				Q1 <b>234</b> ± 25	10.1 (9.8, 10.4)			
				Q2 <b>278</b> ± 9	10.1 (9.8, 10.4)			
					Geometric mean (95% CI)			
					24.5 (23.8, 25.3)			
					25.6 (24.8, 26.4)			

Author	Type of study	Subjects n, sex, age, country	*Intervention/ design (trials) *Intake measurement (cross-sectional studies)	*Duration (trials) *Choline intake (mg/day) (cross-sectional studies)	tHcy in plasma (µmol/L)		Comment on tHcy in plasma	Other outcomes
					Plasma tHcy (fasting)	Post-methionine (0.1 g/kg bw)		
Chiuvè et al. (2007)	Cross-sectional study within long-term cohort; Nurses' Health Study (NHS) and NHS 2; start 1976 and 1989, respectively	1,477 (healthy premenopausal women), 867 NHS (30–55 years at inclusion), 510 NHS2 (25–42 years at inclusion), USA	FFQ 1984, 1986, 1990 for NHS, and 1991, 1995, 1999 for NHS 2	Q3 <b>305</b> ± 7 Q4 <b>334</b> ± 10 Q5 <b>379</b> ± 36  Energy-adjusted intake Total choline (all forms)  Quintiles (median) Q1 <b>265</b> Q2 <b>297</b> Q3 <b>323</b> Q4 <b>345</b> Q5 <b>385</b>	9.7 (9.5, 10.0) 9.7 (9.4, 9.9) 9.7 (9.4, 9.9) p for trend 0.001  Adjusted for age  Median ± SEM 11.4 ± 0.3 10.7 ± 0.2 10.7 ± 0.2 10.2 ± 0.2 10.1 ± 0.3	24.0 (23.3, 24.8) 24.4 (23.7, 25.2) 24.3 (23.6, 25.0) N.S.	strongest for GPC and stronger for men than women  Inverse relationship between plasma tHcy (age-adjusted, or further adjusted for diet and other lifestyle factors) and (1) total choline intake or (2) choline intake from PChol and GPC, particularly if folate intake is low; no relationship when further adjusting for riboflavin and folate intake	

ALT: alanine transaminase; AST: aspartate transaminase; bw: body weight; CI: confidence interval; DMG: dimethylglycine; FFQ: food frequency questionnaire; GGT:  $\gamma$ -glutamyltransferase; GPC: glycerophosphocholine; HDL: high-density lipoproteins; LDL: low-density lipoproteins; MTHFR: methylene-tetrahydrofolate reductase; n: number of subjects; NHS: Nurses' Health Study; NL: Netherlands; N.S.: not significant; PC: phosphatidylcholine; Q: quintile; RCT: randomised controlled trial; SD: standard deviation; SEM: standard error of the mean; TAG: triacylglycerols; tHcy: total homocysteine.

## Appendix C – SNPs of genes coding for enzymes involved in choline metabolism and their impact on choline requirement and/or risk to develop organ dysfunction while being fed a low-choline diet

Enzyme gene	rs number	Base pair and change	Comments
Phosphatidylethanolamine methyltransferase ( <i>PEMT</i> ) (about 100 SNPs)	rs12325817 rs4646343 rs37601188	–744G→C C→A G→A	<p>Three SNPs that decrease the oestrogen responsive <i>PEMT</i> induction (da Costa et al., 2014) associated with increased risk of choline deficiency on choline depletion. May increase the dietary requirement of choline. Eighteen of 23 female carriers of the variant rs12325817 allele developed organ dysfunction on choline depletion (OR of 25; 95% CI 2.0–256.0; <math>p = 0.002</math>), but men did not (da Costa et al., 2006a). Fischer et al. (2010a) found a gene dose–response relationship in 27 premenopausal women to develop signs of choline deficiency on choline depletion: 80%, 43% and 13% with two, one and zero variant alleles, respectively, developed liver dysfunction. Eleven of 22 postmenopausal women subjected to the standard choline depletion/repletion experiment who received oestrogen were four times less likely to develop choline-deficiency associated liver dysfunction than 11 women who received placebo</p> <p>The rs12325817 CC genotype was associated with an increased risk for breast cancer mortality compared to the GG genotype (OR 1.30, 95% CI 1.01–1.67) (Xu et al., 2008)</p> <p>About 75% of the North Carolina population is carrier of at least one rs12325817 C allele and 18% are homozygous for the variant allele (Corbin and Zeisel, 2012). The rs12325817 allele was associated in 92% of 64 women with a rs4646343 allele (Kohlmeier et al., 2005; da Costa et al., 2006a; Resseguie et al., 2011)</p>
	rs7946(3)	+5465G→A	<p>Despite 30% loss of function, no increased susceptibility to choline deficiency (da Costa et al., 2006a). The <i>PEMT</i> rs79463 SNP is found more frequently in people with fatty liver consuming a low-choline diet (Ivanov et al., 2009) and in 67.9% of patients with NAFLD (healthy subjects 40.7%) (Song et al., 2005)</p> <p>Attenuates rise in plasma Hcy in men with the <i>MTHFR</i> 677TT genotype (Caudill et al., 2009)</p> <p>Gene frequency in 43 Mexican-American women: GG 3, GA 19, AA 21 (Ivanov et al., 2009)</p> <p>All effect alleles of <i>PEMT</i> occur frequently in American subjects of European origin (homozygosity 24–60%), followed by Mexican origin (9–12%) and least in subjects of Asian or African descent (da Costa et al., 2014)</p> <p>Carrier status of offspring without effect on umbilical cord blood choline and its metabolites (Visentin et al., 2015)</p>
Methylenetetrahydrofolate dehydrogenase 1 ( <i>MTHFD1</i> )	rs2236225	1958G→A–	<p>Decreases the availability of methyl-THF for Hcy remethylation and increases reliance on choline-derived methyl groups. May increase the dietary requirement of choline and reduce the synthesis of PC (Ivanov et al., 2009): in a choline depletion/repletion study on 54 healthy adults (<math>n = 26</math> men and <math>n = 28</math> women), more than half of the participants developed organ dysfunction associated with choline deficiency. Signs of choline deficiency were significantly (<math>&gt; 15</math> times in premenopausal women) more likely to occur in subjects</p>



Enzyme gene	rs number	Base pair and change	Comments
			<p>who were carriers of the A allele of the SNP rs2236225 of <i>MTHFD1</i> (OR 7.0; 95% CI 2.0–25.0, <math>p &lt; 0.01</math>) than in non-carriers during the low-choline diet, unless they were also treated with a folic acid supplement (Kohlmeier et al., 2005)</p> <p>Homozygous mothers for the SNP were found to have a 1.5–2-fold increased risk of carrying a child with an NTD (Brody et al., 2002)</p> <p>Carrier status of offspring without effect on umbilical cord blood choline and its metabolites (Visentin et al., 2015)</p> <p>63% of subjects investigated in North Carolina possessed at least one allele of this SNP and 11% were homozygous carriers (da Costa et al., 2006a; Corbin and Zeisel, 2012)</p>
Choline dehydrogenase ( <i>CHDH</i> )	rs9001	+114A→C	<p>Carriers may be protected against organ dysfunction upon choline depletion (OR 0.2; 95% CI 0.05–0.7, <math>p = 0.03</math>) (da Costa et al., 2006a)</p>
	rs12676	+233G→T	<p>May increase the dietary requirement of choline in carriers of the variant associated with increased susceptibility to choline deficiency upon choline depletion in premenopausal women (OR 20.0; 95% CI 1.0–282.0; <math>p = 0.04</math>) (da Costa et al., 2006a)</p> <p>The T allele was associated with an increased risk (OR 1.19, 95% CI 1.00–1.41) for breast cancer compared to the major G allele (Xu et al., 2008)</p> <p>Forty and 75% lower ATP concentration in sperm of men with GT (<math>n = 18</math>) and TT (<math>n = 5</math>) genotypes compared to the GG (<math>n = 17</math>) genotype, respectively (Johnson et al., 2012). The TT genotype is present in 9% of the North Carolina population, the prevalence of the GT genotype is 45% (Johnson et al., 2012)</p> <p>Carrier status of offspring without effect on umbilical cord blood choline and its metabolites (Visentin et al., 2015)</p>
Betainehomocysteine methyltransferase ( <i>BHMT</i> )	rs3733890	+742G→A	<p>Not associated with susceptibility to choline deficiency (da Costa et al., 2006a)</p> <p>This polymorphism was not associated with breast cancer risk (Xu et al., 2008), but with a reduced risk of breast cancer mortality (Xu et al., 2009). Carrier status of offspring without effect on umbilical cord blood choline and its metabolites (Visentin et al., 2015)</p>
Choline kinase A ( <i>CHKA</i> )	rs7928739 rs10791957 rs2512612	A→C A→C A→G	<p>Three SNPs associated with a decreased risk for organ dysfunction on choline depletion in homozygotes (da Costa et al., 2014)</p> <p>Frequency is highest in subjects of African descent followed by Asian and European origin and least frequent in subjects of Mexican origin (da Costa et al., 2014)</p> <p>In a case–control study on 103 cases of spina bifida and of 338 controls, the CHK SNP (rs7928739) genotype with at least one C allele was associated with a reduced risk of spina bifida (OR = 0.60, 95% CI = 0.38–0.94 (Enaw et al., 2006)</p>
	rs6591331	A→T	<p>Associated with increased risk for organ dysfunction in postmenopausal women on choline depletion in homozygotes (da Costa et al., 2014)</p>

Enzyme gene	rs number	Base pair and change	Comments
Choline kinase B ( <i>CHKB</i> )	rs1557502	G→A	Associated with an increased risk for muscle damage on choline depletion (da Costa et al., 2014). Most frequent in subjects of African descent, least frequent with European origin (da Costa et al., 2014). Nine of ten subjects who developed muscle damage were heterozygous or homozygous carriers of the effect alleles for <i>SLC44A1</i> rs2771040 (G) and <i>CHKB</i> rs1557502 (A)
CTP:phosphocholine cytidyltransferase ( <i>CCT</i> )	rs939883	T→A	In a case-control study on 103 cases of spina bifida and of 338 controls, the <i>CCT</i> rs939883 genotype AA was associated with an increased risk of spina bifida (OR = 1.89, 95% CI = 0.97–3.67) (Enaw et al., 2006)
Solute carrier 44A1 (choline transporter) ( <i>SLC44A1</i> )	rs7873937 rs2771040 rs6479313 rs16924529 rs3199966	C→G A→G C→G G→A A→C	Associated with an increased risk for muscle damage on choline depletion with a low-choline diet (da Costa et al., 2014). Nine of ten subjects who developed muscle damage were heterozygous or homozygous carriers of the effect alleles for <i>SLC44A1</i> rs2771040 (G) and <i>CHKB</i> rs1557502 (A) Most frequent in subjects of African descent, least frequent with Asian origin (da Costa et al., 2014)
Methylenetetrahydrofolate reductase ( <i>MTHFR</i> )	rs1801133	677C→T	Thermolabile enzyme, increases the reliance on choline-derived methyl groups for Hcy remethylation when folate intake is insufficient (Yan et al., 2011). Significantly increased plasma Hcy, decreased plasma PC and SPM with low folate status/intake in both men and women with either CC (n = 28) or TT (n = 17) genotype, but no change in plasma choline and leukocyte global DNA methylation. Women with the TT genotype had a 10.3% increase in plasma PC while consuming adequate amounts of folate and choline. No changes in plasma PC in response to diet in subjects with the CC genotype (Abratte et al., 2009) In 60 healthy men, 29 with the TT genotype and 31 with the CC genotype, an intake of 300 mg choline/day for 12 weeks was sufficient to maintain liver and kidney function, but 438 µg DFE/day did not prevent a rise in plasma tHcy in subjects with the TT genotype. Under these conditions, choline supplementation (up to 1,900 mg/day) had no effect on plasma tHcy and serum folate concentrations. Choline intake decreased DNA methylation in subjects with the CC genotype but not in TT subjects (Solis et al., 2008; Veenema et al., 2008; Caudill et al., 2009) Carrier status of offspring without effect on umbilical cord blood choline and its metabolites (Visentin et al., 2015) TT genotype frequency varies between ethnic groups (2–35%)
	rs1801131	1298A→C	Reduced enzyme activity; no association with risk for choline deficiency in choline depletion/repletion studies (Kohlmeier et al., 2005)

ATP: adenosine triphosphate; BHMT: betaine-homocysteine methyltransferase; CCT/CTP:phosphocholine cytidyltransferase; CHDH: choline dehydrogenase; CHK: choline kinase; CI: confidence interval; DFE: dietary folate equivalent; DNA: deoxyribonucleic acid; Hcy: homocysteine; MTHFD1: 5,10-methylene-tetrahydrofolate dehydrogenase 1; MTHFR: methylene-tetrahydrofolate reductase; NAFLD: non-alcoholic fatty liver disease; NTD: neural tube defect; OR: odds ratio; PC: phosphatidylcholine; PEMT: phosphatidylethanolamine N-methyltransferase; rs number: Reference SNP cluster ID; SLC44A1: solute carrier family 44 member 1 (choline transporter); SNP: single-nucleotide polymorphism; SPM: sphingomyelin; THF: tetrahydrofolate.

## Appendix D – Depletion/repletion studies for choline

(choline intake per 70-kg body weight per day)

Author	Aim of investigation	Outcome measurements	Participants	Design/duration	Results	Comment
Zeisel et al. (1991) <b>1*</b>	Experimental choline deficiency in humans	Choline, PC in plasma; PC in red blood cells; liver and kidney function, blood lipids, liver size and density by CT	Male, n = 15, healthy <b>A controls</b> n = 6, mean age 26.8 years; <b>B depleted</b> n = 8, mean age 29.1 years One recruited control subject was excluded (abnormal liver function tests on day 1)	Metabolic unit; <b>Week 1:</b> A and B: baseline diet (13 mg/70-kg bw per day) + 500 mg/day choline <b>Week 2-4:</b> <b>A:</b> baseline diet + 500 mg/day choline <b>B:</b> baseline diet + placebo	<b>Week 1:</b> free choline in plasma 9.6–10.9 $\mu$ mol/L; plasma PC 1.3–2.0 mmol/L <b>Week 4:</b> <b>A:</b> no change in plasma choline/PC, increase by 14% in red blood cell PC, no change in ALT <b>B:</b> choline in plasma decreased by 30%, plasma PC (as % of day 7 value) decreased by 30%, decrease in red blood cell PC by 15%; significant <b>increase in ALT</b> by 50%; non-significant increase in liver size <b>Week 5:</b> <b>A:</b> no change <b>B:</b> plasma choline, plasma PC, ALT return to baseline	Plasma choline, plasma PC and serum ALT activity expressed as a change from day 7 to day 28 Three-week depletion of dietary choline (513–13 mg choline/day) significantly decreased plasma choline and PC and increases serum ALT activity in all subjects. No effects on other hepatic or kidney function parameters
Kohlmeier et al. (2005) <b>2</b>	Influence of genetic variants of folate metabolism on susceptibility to choline deficiency	Liver fat by MRI, CK in serum, Plasma folate, plasma tHcy, SAM, SAH; tHcy response to methionine load before and after depletion; genotyping for <i>MTHFR</i> , <i>MTHFD1</i> and <i>RFC1</i>	n = 54, female n = 28, mean age 38.7 years, healthy	<b>Week 5</b> (i.e. 35 days): <b>A:</b> baseline diet + 500 mg/day choline <b>B:</b> baseline diet + 500 mg/day choline Metabolic unit <b>Baseline:</b> 10 days, 550 mg choline/70-kg bw per day + 400 $\mu$ g folic acid <b>Depletion</b> (up to 42 days): < 50 mg choline/70-kg bw per day and 100 $\mu$ g folate/day	<b>Organ dysfunction</b> 12/54 subjects 5-fold increase in CK 24/54 increase (at least by 28%) in liver fat content, no effect of folate intake <b>Genotyping and %</b>	More than 50% of the participants developed signs of organ dysfunction when consuming < 50 mg/70-kg bw per day. Susceptibility to develop signs of choline deficiency on a 50 mg/70-kg bw per

Author	Aim of investigation	Outcome measurements	Participants	Design/duration	Results	Comment
da Costa et al. (2005) <b>3</b>	Choline deficiency and capacity to methylate tHcy	Total plasma tHcy, before and after Met load (100 mg/kg bw) before and after choline depletion and repletion, plasma choline, betaine, PC, folate liver fat by MRI	n = 8 males, age 20-46 years, healthy	<p><b>A</b> plus 400 µg folic acid/day</p> <p><b>B</b> placebo</p> <p><b>Repletion</b> (increasing amount (137–550 mg/70-kg bw per day) up to &gt; 550 mg choline per day for ≥ 3 days)</p>	<p><b>symptomatic choline deficiency:</b></p> <p><i>MTHFD1</i> 1958 GG n = 20: 40%</p> <p><i>MTHFD1</i> 1958 GA n = 28: 82%</p> <p><i>MTHFD1</i> 1958 AA n = 6: 83%</p> <p>GG vs GA/AA <b>OR 7.0 (95% CI 2.0–25)</b></p> <p><b>p = 0.007</b></p> <p><i>RFC1</i> 80 AG n = 20: 70%</p> <p><i>RFC1</i> 80 GG n = 15: 73%</p> <p>AA vs AG/GG OR 1.82 (95% CI 0.56-5.9) N.S</p> <p>Mean serum folate significantly lower in subjects with low folate intake (22.1 (B) vs 28.3 mmol/L (A)) without effect by genetic polymorphism</p>	<p>day-choline diet greater in carriers of the <i>MTHFD1</i> G1958A polymorphism: OR 7.0 (95% CI 2.0–25; p &lt; 0.01) unless they received additional folic acid</p> <p>Susceptibility to develop signs of choline deficiency not influenced by polymorphism of <i>MTHFR</i> or <i>RFC1</i></p>
				<p>Standardised depletion/repletion design</p> <p><b>Baseline</b> diet (10 days): 550 mg choline/70-kg bw per day + 400 DFE/day</p> <p><b>Depletion</b> diet (up to 42 days): &lt; 50 mg choline/70-kg bw per day</p> <p><b>Repletion</b> diet:</p> <p>1) Subjects not clinically choline deficient: 550 mg choline diet for 3 days</p> <p>2) Subjects clinically choline deficient: graded amounts</p>	<p><b>Organ dysfunction</b></p> <p>4/8 increase in liver fat</p> <p><b>tHcy in plasma</b></p> <p><b>Depletion</b> fasting tHcy significantly increased by 1.3 µmol/L in clinically choline-deficient participants (no significant change in the non-deficient subjects)</p> <p><b>Free choline in plasma (mean)</b></p> <p>Before depletion 10 µmol/L</p>	<p>Half of the participants developed signs of liver dysfunction when consuming &lt; 50 mg choline/70-kg bw per day; no difference in change in plasma choline (or betaine) between those with and without organ dysfunction</p>

Author	Aim of investigation	Outcome measurements	Participants	Design/duration	Results	Comment
da Costa et al. (2006a) <b>3</b>	Choline deficiency and lymphocyte apoptosis and DNA damage	CK, liver fat by MRI, 24 h urine choline and betaine, plasma folate, peripheral lymphocytes at baseline, after depletion and repletion: DNA fragmentation (TUNEL) and strand breaks (COMET), activated caspase-3 (used as a marker for apoptosis)	n = 51, n = 31 female, age 18-70 years, healthy	of choline sequentially in 10 days periods (138, 275, 413, 550 mg/70-kg bw per day until hepatic steatosis resolved)	Clinically depleted 7 $\mu\text{mol/L}$ Not clinically depleted 7 $\mu\text{mol/L}$ <b>PC in plasma (mean)</b> Before depletion 1,818 $\mu\text{mol/L}$ Clinically depleted 1,564 $\mu\text{mol/L}$ Not clinically depleted 1,834 $\mu\text{mol/L}$ <b>Betaine in plasma (mean)</b> Before depletion 66 $\mu\text{mol/L}$ Clinically depleted 36 $\mu\text{mol/L}$ Not clinically depleted 34 $\mu\text{mol/L}$	Choline deficiency is associated with <i>in vitro</i> signs of DNA damage and of apoptosis in peripheral lymphocytes
				Metabolic unit Standardised depletion/repletion design <b>Baseline diet</b> (10 days): 550 mg choline/70-kg bw per day + 400 DFE/day <b>Depletion diet</b> (up to 42 days): < 50 mg choline/70-kg bw per day and 100 DFE/day <b>A</b> plus 400 $\mu\text{g}$ folic acid/day, n = 26 <b>B</b> placebo, n = 25 <b>Repletion diet</b> 1) Subjects not clinically choline deficient: 550 mg choline diet for 3 days 2) Subjects clinically choline	<b>Organ dysfunction</b> 33/51, including 26/51 liver dysfunction (18 females) 1/51 muscle dysfunction only 6/51 both liver and muscle dysfunction returning to normal after choline repletion <b>Plasma folate</b> Significant decrease during choline depletion without extra folic acid: 26.0–21.4 $\mu\text{mol/L}$ (and p = 0.0003 without folate supplementation) <b>24 h urine choline</b>	



Author	Aim of investigation	Outcome measurements	Participants	Design/duration	Results	Comment
Fischer et al. (2007) <b>3</b>	Dietary requirement in healthy men and women and clinical sequelae of choline deficiency	Plasma choline, PC, SAM, SAH, Met, tHcy, methylglycine and DMG CK, Liver fat by MRI	n = 57, n = 16 premenopausal women, n = 15 postmenopausal women, n = 26 men Age 18-70 years, healthy	deficient: graded amounts of choline sequentially in 10 days periods (137.5, 275, 412.5 and 550 mg/70-kg bw per day and > 550 mg for 3 days)	<b>and betaine</b> Decrease from about 25 to 10 and from 80 to about 30 $\mu\text{mol/g}$ creatinine, respectively, with choline depletion <b>Activated caspase-3 assay in lymphocytes</b> Higher amounts in cells from clinically choline deficient subjects, compared to non-deficient subjects ( $p < 0.05$ ) <b>TUNEL assay</b> More TUNEL-positive lymphocyte cells during choline depletion with or without organ dysfunction, without folic acid supplement ( $p = 0.026$ ) <b>COMET assay</b> COMET-Tail moment increase during choline depletion compared to baseline	Most men and postmenopausal women (68.4%) developed clinical choline deficiency when on < 50 mg choline/day independent on folate intake 18/57 subjects did not develop signs of choline
				Metabolic unit Standardised depletion/repletion design <b>Baseline</b> diet (10 days): 550 mg choline/70-kg bw per day + 400 DFE/day <b>Depletion</b> diet (up to 42 days): < 50 mg choline/70-kg bw per day + 100 DFE/day	1) <b>6 while on 550 mg choline baseline diet (550 mg/70 kg bw per day)</b> , all men 2) <b>33 while on</b>	

Author	Aim of investigation	Outcome measurements	Participants	Design/duration	Results	Comment
				<p><b>A</b> plus 400 µg folic acid/day</p> <p><b>B</b> placebo</p> <p><b>Repletion diet:</b></p> <p>1) Subjects not clinically choline deficient: 550 mg choline diet for 3 days</p> <p>2) Subjects clinically choline deficient: graded amounts of choline sequentially in 10 days periods (137.5, 275, 412.5 and 550 mg/70-kg bw per day, then &gt; 550 mg for 3 days)</p>	<p><b>low-choline diet (50 mg/70 kg bw per day):</b> 14/20 men (70%), 7/16 (44%) premenopausal women</p> <p>12/15 (80%) postmenopausal women; with <b>liver steatosis</b> alone: in 8/20 men, 12/15 postmenopausal women and 6/16 premenopausal women</p> <p><b>Choline (metabolites) in plasma on depletion:</b></p> <p>Choline decrease by 28–33%, betaine by ≈50%, PC only in subjects with organ dysfunction, Met decreased only in subjects with organ dysfunction, DMG and MG decreased, tHcy increased, SAM and SAH did not change</p> <p>Serum uric acid increased in all subjects during depletion</p> <p><b>Repletion of choline depleted subjects: see Table 2, Section 5.1.1.3</b></p>	<p>deficiency with &lt; 50 mg choline/day</p>

Author	Aim of investigation	Outcome measurements	Participants	Design/duration	Results	Comment
Niculescu et al. (2007) 3	Organ dysfunction on low-choline diet and SNPs in genes involved in choline and folate metabolism	Liver fat by MRI, CK in serum, Peripheral lymphocytes at 10 days and after depletion for genotyping <i>MTHFD1</i> , <i>PEMT</i> , <i>CHDH</i> and for change in expression with low-choline diet and DNA methylation	n = 33, age 20-67 years, 19 women, healthy	Metabolic unit Standardised depletion/repletion design <b>Baseline</b> diet (10 days): 550 mg choline/70-kg bw/day + 400 DFE/day <b>Depletion</b> diet (up to 42 days): < 50 mg choline/70-kg bw per day <b>A</b> plus 400 µg folic acid/day <b>B</b> placebo <b>Repletion</b> diet	No outcome measurements indicative of choline requirement	Previous studies showed that the <i>PEMT</i> (rs12325817) and <i>MTHFD1</i> (rs2236225) SNPs predispose subjects to develop organ dysfunction when they consume a low-choline diet (Kohlmeier et al., 2005; da Costa et al., 2006a) At baseline, subjects with the <i>PEMT</i> (rs12325817) and <i>MTHFD1</i> (rs2236225) SNPs, compared with subjects without the SNPs, had a different expression of genes involved in apoptosis, the DNA damage checkpoint, and cell proliferation control This suggests that the presence of the <i>PEMT</i> and <i>MTHFD1</i> genotypes can lead to differences in the phenotypes at baseline (i.e. even before consuming a low-choline diet). Subjects may differ in their susceptibility to dietary choline deficiency. In women who are carriers of the <i>PEMT</i> allele, the risk of choline deficiency is higher

Author	Aim of investigation	Outcome measurements	Participants	Design/duration	Results	Comment
Fischer et al. (2010a) 4	Low-choline related organ dysfunction, in relation to number of alleles of rs12325817 in premenopausal women, and in relation to oestrogen in postmenopausal women	Liver fat by MRI, CK, AST, ALT Plasma choline (metabolites) Genotyping for <i>PEMT</i> rs12325817	<b>A:</b> n = 27 premenopausal women, age 18–49 years. <b>B:</b> n = 22 postmenopausal women, age 50–73 years, randomised to receive oestrogen ( <b>B1</b> ) or placebo ( <b>B2</b> ). Healthy	Metabolic unit Standardised depletion/repletion <b>Baseline diet</b> (10 days): 550 mg choline/70-kg bw per day <b>Depletion diet</b> (up to 42 days): < 50 mg choline/70-kg bw per day <b>Repletion diet:</b> 550–850 mg/70-kg bw per day for up to 10 days. If signs of organ dysfunction did not resolve after 10 days of repletion diet: <i>ad libitum</i> diet for 2 weeks	<b>Among premenopausal women:</b> 11/27 developed choline deficiency/organ dysfunction. There was a dose-response effect of rs12325817 on the risk of choline related organ dysfunction: 80%, 43%, and 13% of women with 2, 1, or 0 alleles, respectively, developed organ dysfunction during the low-choline diet <b>Among postmenopausal women:</b> Only 2/11 (18%) who received oestrogen ( <b>B1</b> ) and 8/11 (73%) who received placebo ( <b>B2</b> ), developed organ dysfunction during the low-choline diet	Dietary requirement for choline is higher in postmenopausal women (because of their lower oestrogen concentrations) than in premenopausal women Choline requirements for both groups of women are further increased by rs12325817. 80% of homozygous women develop organ dysfunction on the depletion diet vs 43% of those with one copy and 13% of women homozygous for the wild-type No oestrogen vs oestrogen increases fourfold the risk for organ dysfunction on the depletion diet. Oestrogen mitigates the effect of the <i>PEMT</i> SNP. Oestrogen may decrease choline requirement in postmenopausal women
Sha et al. (2010) 3	Metabolomic profiling to predict organ dysfunction with deficient choline intake	Liver fat by MRI CK, AST, ALT Plasma choline (metabolites), Met, Hcy, sarcosine, DMG, cysteine, cystathionine, Metabolomic analysis of plasma	n = 53, n = 30 women, age 18–70 years, healthy	Metabolic unit Standardised depletion/repletion design <b>Baseline diet</b> (10 days): 550 mg choline/70-kg bw per day <b>Depletion diet</b> (up to 42 days): < 50 mg choline/70-kg bw per day <b>Repletion diet</b> ( $\geq$ 3 days,	<b>Organ dysfunction</b> <b>Baseline diet:</b> 9 (17%) developed fatty liver (n = 4) or muscle dysfunction (n = 5), without special metabolome <b>Depletion</b> (n = 44): 23 fatty liver, 5 muscle dysfunction	Metabolomic profiles of subjects at baseline could predict the development of liver dysfunction when deprived of dietary choline

Author	Aim of investigation	Outcome measurements	Participants	Design/duration	Results	Comment
Spencer et al. (2011) <b>4</b>	Choline deficiency and hepatic steatosis and gut microbiome	Liver fat by MRI CK AST, ALT Sequencing of the 16S RNA bacterial genes in stool; genotyping of <i>PEMT</i> promoter SNP rs12325817	n = 15 females, age not reported, healthy	(≥ 550 mg/70-kg bw per day)	Higher plasma Hcy, cysteine, cystathionine, keto-acids at baseline in subjects who later develop fatty liver Choline deficiency increased plasma carnitine and acyl-carnitine, decreased pyridoxate Baseline plasma choline has no predictive value	Host factors as well as gut bacteria respond to dietary choline deficiency, but individual microbiota persist although all subjects consumed the same diets
da Costa et al. (2011) <b>3+4</b>	PC-DHA plasma concentration used as a non-invasive marker of liver <i>PEMT</i> activity	Plasma DHA, PC-DHA, ratio PC-DHA/total PC	n = 72, age 18-70 years; n = 20 men; n = 52 women of which n = 25 postmenopausal and n = 27 premenopausal	Standardised depletion/repletion design 2 months: <b>Baseline</b> diet (10 days): 550 mg choline/70-kg bw per day <b>Depletion</b> diet (up to 42 days): < 50 mg choline/70-kg bw per day <b>Repletion</b> diet (10 days, (≥ 850 mg/70-kg bw per day)	70% of the subjects possess at least one <i>PEMT</i> rs12325817 allele	Plasma ratio PC-DHA/total PC higher in premenopausal women than men or postmenopausal (at baseline and even when on a low-choline diet) Plasma PC-DHA/total PC at baseline and <i>PEMT</i> activity in liver: lower in premenopausal women homozygous for the rs12325817 polymorphism in the <i>PEMT</i> gene



Author	Aim of investigation	Outcome measurements	Participants	Design/duration	Results	Comment
da Costa et al. (2011) <b>3+4</b>	Identification of effect alleles of SNPs known to influence dietary requirement for choline	DNA concentration by spectrometry; genotyping of alleles	n = 79, 18-70 years old; n = 26 men n = 53 women of which n = 26 post premenopausal and n = 27 premenopausal	Standardised depletion/repletion design <b>Baseline</b> diet (10 days): 550 mg choline/70-kg bw per day <b>Depletion</b> diet (up to 42 days): < 50 mg choline/70-kg bw per day <b>Repletion</b> diet	Effect alleles identified of SNPs in genes for the choline transporter (SLC44A1) and choline kinase A and B (see Appendix C) Choline deficiency related organ dysfunction (liver or muscle: 50/79, including 20 of 26 postmenopausal women, 11 of 27 premenopausal women 19 of 26 men	29 of 79 healthy subjects did not develop organ dysfunction while consuming a low-choline diet for 6 weeks

\*Same numbers in the column "author" indicate references providing data from the same cohort.

ALT: alanine aminotransferase; AST: aspartate aminotransferase; CHDH: choline dehydrogenase; CI: confidence interval; CK: creatine kinase; COMET: single-cell gel electrophoresis; CT: computerised tomography; DFE: dietary folate equivalent; DHA: docosahexaenoic acid; DMG: dimethylglycine; DNA: deoxyribonucleic acid; LDH: lactate dehydrogenase; Met: methionine; MG: methylglycine; MRI: magnetic resonance imaging; MTHFD1: 5,10-methylene-tetrahydrofolate dehydrogenase 1; MTHFR: Methylene-tetrahydrofolate reductase; N.S.: not significant; OR: odds ratio; PC: phosphatidylcholine; PEMT: phosphatidylethanolamine N-methyltransferase; RFC1: reduced folate carrier 1; RNA: ribonucleic acid; rs number: Reference SNP cluster ID; SAH: S-adenosylhomocysteine; SAM: S-adenosyl-methionine; SLC44A1: solute carrier family 44 member 1 (choline transporter); SNP: single-nucleotide polymorphism; tHcy: total homocysteine; TUNEL: terminal deoxynucleotidyl transferase mediated dUTP nick end labelling.